



Beachwatch

State of the beaches 2021–22

Sydney Region

© 2022 State of NSW and Department of Planning and Environment (DPE)

With the exception of photographs, the State of NSW and Department of Planning and Environment are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required for the reproduction of photographs.

The Department of Planning and Environment (DPE) has compiled this report in good faith, exercising all due care and attention. No representation is made about the accuracy, completeness or suitability of the information in this publication for any particular purpose. DPE shall not be liable for any damage which may occur to any person or organisation taking action or not on the basis of this publication. Readers should seek appropriate advice when applying the information to their specific needs.

Every effort has been made to ensure that the information in this document is accurate at the time of publication. However, as appropriate, readers should obtain independent advice before making any decision based on this information.

All content in this publication is owned by DPE and is protected by Crown Copyright, unless credited otherwise. It is licensed under the Creative Commons Attribution 4.0 International (CC BY 4.0), subject to the exemptions contained in the licence. The legal code for the licence is available at Creative Commons.

DPE asserts the right to be attributed as author of the original material in the following manner: © State of New South Wales and Department of Planning and Environment 2022.

Front cover: Shelly Beach, Cronulla (Cameron Board/DPE)

Published by:

Environment and Heritage

Department of Planning and Environment

Locked Bag 5022, Parramatta NSW 2124

Phone: +61 2 9995 5000 (switchboard)

Phone: 1300 361 967 (Environment, Energy and Science enquiries)

TTY users: phone 133 677, then ask for 1300 361 967

Speak and listen users: phone 1300 555 727, then ask for 1300 361 967

Email: info@environment.nsw.gov.au

Website: www.environment.nsw.gov.au

Report pollution and environmental incidents

Environment Line: 131 555 (NSW only)

or info@environment.nsw.gov.au

See also www.environment.nsw.gov.au

ISBN 978-1-922840-72-1

EHG2022/0322

July 2022

Contents

Sydney region summary 2021–2022	1
Beach monitoring in NSW	1
Rainfall impacts	1
Northern Sydney (Pittwater to Manly)	10
Overall results	10
Management	13
Palm Beach	17
Whale Beach	18
Avalon Beach	19
Bilgola Beach	20
Newport Beach	21
Bungan Beach	22
Mona Vale Beach	23
Warriewood Beach	24
Turimetta Beach	25
North Narrabeen Beach	26
Narrabeen Lagoon (Birdwood Park)	27
Bilarong Reserve	28
Collaroy Beach	29
Long Reef Beach	30
Dee Why Beach	31
North Curl Curl Beach	32
South Curl Curl Beach	33
Freshwater Beach	34
Queenscliff Beach	35
North Steyne Beach	36
South Steyne Beach	37
Shelly Beach	38
Barrenjoey Beach	39
Paradise Beach Baths	40
Clareville Beach	41

Taylors Point Baths	42
Bayview Baths	43
Elvina Bay	44
North Scotland Island	45
South Scotland Island	46
The Basin	47
Great Mackerel Beach	48
Central Sydney (Bondi to Little Bay & Sydney Harbour)	49
Overall results	49
Management	52
Bondi Beach	65
Tamarama Beach	66
Bronte Beach	67
Clovelly Beach	68
Gordons Bay	69
Coogee Beach	70
Maroubra Beach	71
South Maroubra Beach	72
South Maroubra Rockpool	73
Malabar Beach	74
Little Bay Beach	75
Camp Cove	76
Watsons Bay	77
Parsley Bay	78
Nielsen Park	79
Rose Bay Beach	80
Murray Rose Pool	81
Dawn Fraser Pool	82
Chiswick Baths	83
Cabarita Beach	84
Woolwich Baths	85
Tambourine Bay	86
Woodford Bay	87
Greenwich Baths	88
Hayes Street Beach	89


Clifton Gardens	90
Balmoral Baths	91
Edwards Beach	92
Chinamans Beach	93
Northbridge Baths	94
Davidson Reserve	95
Gurney Crescent Baths	96
Clontarf Pool	97
Forty Baskets Pool	98
Fairlight Beach	99
Manly Cove	100
Little Manly Cove	101
Southern Sydney (Sutherland beaches, lower Georges River, Botany Bay & Port Hacking)	102
Overall results	102
Management	105
Boat Harbour	113
Greenhills Beach	114
Wanda Beach	115
Elouera Beach	116
North Cronulla Beach	117
South Cronulla Beach	118
Shelly Beach	119
Oak Park	120
Silver Beach	121
Como Baths	122
Jew Fish Bay Baths	123
Oatley Bay Baths	124
Carss Point Baths	125
Sandringham Baths	126
Dolls Point Baths	127
Ramsgate Baths	128
Monterey Baths	129
Brighton-Le-Sands Baths	130
Kyeemagh Baths	131

Foreshores Beach	132
Yarra Bay	133
Frenchmans Bay	134
Congwong Bay	135
Jibbon Beach	136
Horderns Beach	137
GyMEA Bay Baths	138
Lilli Pilli Baths	139
Gunnamatta Bay Baths	140
How to read this report	141
Beach Suitability Grades	141
Explanation of tables	145
Explanation of graphs, charts, and information bars on beach pages	145
References	149
More information	149

Recreational water quality has been monitored in the Sydney region since 1989 by the Department of Planning and Environment's Beachwatch program. This report summarises the performance of 97 swimming sites in the Sydney region, providing a long-term assessment of how suitable a site is for swimming. Monitored sites include ocean beaches, a lagoon, a rockpool and estuarine sites in Pittwater, Sydney Harbour, Botany Bay, lower Georges River and Port Hacking.

In 2021–2022, 81% of swimming sites in the Sydney region were graded as Good or Very Good, including 37 ocean beaches. These sites were suitable for swimming for most or almost all of the time. While this is a decline in overall performance from the previous year, it reflects the extreme wet weather conditions and flooding experienced over summer and early autumn. The Sydney region has a large proportion of lagoon and estuarine swimming locations, which have been most susceptible to impacts from significant rain events.

Sydney region summary 2021–2022



Dee Why Beach

Photo: Beachwatch/DPE

Beach monitoring in NSW

The water quality of beaches and other swimming locations is monitored under the NSW Government's Beachwatch programs to provide the community with accurate information on the cleanliness of the water and to enable individuals to make informed decisions about where and when to swim. Routine assessment also measures the impact of pollution sources, enables the effectiveness of stormwater and wastewater management practices to be assessed and highlights areas where further work is needed.

Swimming sites in NSW are graded as Very Good, Good, Fair, Poor or Very Poor in accordance with the National Health and Medical Research Council's 2008 *Guidelines for Managing Risks in Recreational Waters*. These Beach Suitability Grades provide a long-term assessment of how suitable a beach is for swimming. The grades are determined from the most recent 100 water quality results (2–4 years' worth of data depending on the sampling frequency) and a risk assessment of potential pollution sources.

See the section on **Quality assurance** in the Statewide Summary for results of the quality assurance program.

Recreational water quality has been monitored in the Sydney region since 1989 by the Department of Planning and Environment's Beachwatch program.

A **quality assurance** program ensures the information collected and reported by Beachwatch is accurate and reliable.

During 2021–2022, 97 swimming sites were monitored including ocean beaches, a lagoon, a rockpool and estuarine sites in Pittwater, Sydney Harbour, Botany Bay, lower Georges River and Port Hacking.

Rainfall impacts

Rainfall is the major driver of pollution to recreational waters, generating stormwater runoff and triggering untreated discharges from the wastewater treatment and transport systems. Changes in rainfall patterns are reflected in beach water quality over time due to variation in the frequency and extent of stormwater and wastewater inputs.

The Beach Suitability Grades for 2021–2022 are based on water quality data collected over the last 2–4 years. Rainfall over this period has been diverse:

- 2018–2019: variable rainfall, with a mix of extended dry weather periods and some very wet months
- 2019–2020: average to below average rainfall, except for some isolated wet weather events and wet February


- 2020–2021: variable rainfall with some very wet months over summer and early autumn
- 2021–2022: varied rainfall, with extreme wet weather conditions over summer and early autumn, and significant flooding.

See the section on **How to read this report** on page 141 for an explanation of the graphs, tables and Beach Suitability Grades.

Sydney's total rainfall in winter 2021 was below the long-term average. August experienced above average rainfall, mostly due to heavy rain over 2 consecutive days in late August.

While monthly rainfall totals in September and October were average to below average, the Sydney region received above average rainfall totals in spring 2021 due to a wet November. Mona Vale recorded more than double the long-term monthly average rainfall for November with 203 mm.

Summer 2021–2022 rainfall totals were above average, with February notably wet. Sydney (Observatory Hill) recorded its wettest summer since 1991–1992 with 655 mm of rain over summer. Collaroy had its highest February rainfall total on record with 419 mm, and Randwick and Rose Bay had their highest total February rainfall since 1990, with 422 mm and 409 mm respectively. Heavy downpours in February resulted in flash flooding across Sydney.



Debris and discoloured water at Queenscliff Beach following extreme wet weather and flooding in March 2022

Photo: Beachwatch/DPE

March 2022 was the wettest March on record in many areas of Sydney. Mona Vale and Collaroy had their highest March rainfall totals on record with 577 mm and 686 mm of rain for the month, respectively. Rose Bay, Randwick and Sydney Airport received their highest March rainfall totals on record, with 588 mm, 563 mm and 474 mm of rain for the month.

The severe wet weather resulted in flash flooding across Sydney and major flooding of coastal waterways, including the Hawkesbury–Nepean. Beachwatch issued an extreme wet weather pollution alert on all Sydney daily beach pollution forecasts during March 2022, advising that stormwater pollution and floodwaters may be impacting swimming sites for an extended period, with lifeguard reports of floating debris and discoloured water continuing after the rain had ceased.


The wet weather continued in April 2022 with well above the long-term monthly average rainfall recorded. A heavy rainfall event was recorded on 7–8 April, with Rose Bay receiving a record high daily rainfall total for April of 167 mm, and Randwick recording 155 mm on the 7th. More than 2 and a half times the long-term monthly average rainfall was recorded at Collaroy, Randwick and Rose Bay with 302 mm, 326 mm and 379 mm, respectively.

The continued wet weather in April triggered more flooding in coastal waterways, including the Georges, Woronora, Nepean and Hawkesbury rivers. Beachwatch issued an extreme wet weather pollution alert until the flooding and stormwater impacts had subsided.

Flooding and water quality

Beachwatch monitoring showed flooding events impacted waterways beyond the flood zones, including Sydney Harbour and Sydney ocean beaches, making microbial water quality unsuitable for swimming. Routine monitoring at coastal swim sites in Sydney detected significantly elevated microbial counts at all 97 monitored swim sites, which posed an increased health risk to bathers. The most affected areas were in estuaries, such as Sydney Harbour, which have a lower level of flushing and took longer to recover from the floodwater events than the ocean beaches. Routine water quality testing showed some sites unsuitable for swimming for up to 4 weeks

As microbial levels returned to normal at swim sites monitored by Beachwatch there was still a large amount of debris or other hazards, such as murky water, which posed a risk to recreational activities.



Marine algal bloom present in the water

Photo: Chad Weston/NPWS, DPE

Marine algal blooms

Water NSW reported several occurrences of marine algal blooms at Sydney beaches in late 2021. Blooms of *Trichodesmium* sp. and *Noctiluca* sp. were reported at Sydney beaches in November and December 2021, and in Sydney Harbour in November 2021. Marine algae advisories were issued on the Beachwatch and Water NSW websites.

The appearance of **marine algae** is sometimes mistaken for **sewage contamination** or **oil slicks**, due to a strong odour and red or brown discolouration in the water caused by the blooms.

As a precaution, direct contact with algae should be avoided as it can cause skin and eye irritations. The marine algal blooms dissipated with changes in tide and wind conditions.

Beachwatch issues daily **beach pollution forecasts** to enable beach goers to make informed decisions about where and when to swim.

Pollution forecasts for the Sydney region can be accessed via the [Beachwatch website](#), [email subscription](#), [Twitter](#) and [Facebook](#).

Health risks
















































Contamination of recreational waters with faecal material from animal and human sources can pose significant health problems to beach users owing to the presence of pathogens (disease-causing micro-organisms) in the faecal material. The most common groups of pathogens found in recreational waters are bacteria, protozoans and viruses.

Exposure to contaminated water can cause gastroenteritis, with symptoms including vomiting, diarrhoea, stomach-ache, nausea, headache and fever. Eye, ear, skin and upper respiratory tract infections can also be contracted when pathogens come into contact with small breaks and tears in the skin or ruptures of the delicate membranes in the ear or nose.

Certain groups of users may be more vulnerable to microbial infection than others. Children, the elderly, people with compromised immune systems, tourists, and people from culturally and linguistically diverse backgrounds are generally most at risk.





Beach Suitability Grades for swimming sites in the Sydney region


Swimming site	Site type	Beach Suitability Grade	Change
Northern Sydney – Ocean beaches			
Palm Beach	Ocean beach	VG	○
Whale Beach	Ocean beach	VG	○
Avalon Beach	Ocean beach	VG	○
Bilgola Beach	Ocean beach	G	↓
Newport Beach	Ocean beach	G	↓
Bungan Beach	Ocean beach	G	↓
Mona Vale Beach	Ocean beach	VG	○
Warriewood Beach	Ocean beach	G	○
Turimetta Beach	Ocean beach	G	○
North Narrabeen Beach	Ocean beach	G	○
Narrabeen Lagoon (Birdwood Park)	Lagoon	P	○
Bilarong Reserve	Lagoon	P	○
Collaroy Beach	Ocean beach	G	○
Long Reef Beach	Ocean beach	G	○
Dee Why Beach	Ocean beach	G	↓
North Curl Curl Beach	Ocean beach	G	○
South Curl Curl Beach	Ocean beach	VG	○
Freshwater Beach	Ocean beach	G	○
Queenscliff Beach	Ocean beach	G	○
North Steyne Beach	Ocean beach	G	○
South Steyne Beach	Ocean beach	G	○
Shelly Beach	Ocean beach	G	○
Northern Sydney – Pittwater			
Barrenjoey Beach	Estuarine	G	○
Paradise Beach Baths	Estuarine	G	○

Swimming site	Site type	Beach Suitability Grade	Change
Northern Sydney – Pittwater (continued)			
Clareville Beach	Estuarine		
Taylor's Point Baths	Estuarine		
Bayview Baths	Estuarine		
Elvina Bay	Estuarine		
North Scotland Island	Estuarine		
South Scotland Island	Estuarine		
The Basin	Estuarine		
Great Mackerel Beach	Estuarine		
Central Sydney – Ocean beaches			
Bondi Beach	Ocean beach		
Tamarama Beach	Ocean beach		
Bronte Beach	Ocean beach		
Clovelly Beach	Ocean beach		
Gordons Bay	Ocean beach		
Coogee Beach	Ocean beach		
Maroubra Beach	Ocean beach		
South Maroubra Beach	Ocean beach		
South Maroubra Rockpool	Ocean baths		
Malabar Beach	Ocean beach		
Little Bay Beach	Ocean beach		
Central Sydney – Sydney Harbour			
Camp Cove	Estuarine		
Watsons Bay	Estuarine		
Parsley Bay	Estuarine		
Nielsen Park	Estuarine		
Rose Bay Beach	Estuarine		

Swimming site	Site type	Beach Suitability Grade	Change
Central Sydney – Sydney Harbour (continued)			
Murray Rose Pool	Estuarine	G	○
Dawn Fraser Pool	Estuarine	G	○
Chiswick Baths	Estuarine	G	○
Cabarita Beach	Estuarine	G	○
Woolwich Baths	Estuarine	G	○
Tambourine Bay	Estuarine	P	○
Woodford Bay	Estuarine	G	○
Greenwich Baths	Estuarine	G	○
Hayes St Beach	Estuarine	G	○
Clifton Gardens	Estuarine	G	○
Balmoral Baths	Estuarine	G	○
Edwards Beach	Estuarine	G	○
Chinamans Beach	Estuarine	G	○
Northbridge Baths	Estuarine	P	↓
Davidson Reserve	Estuarine	P	○
Gurney Crescent Baths	Estuarine	F	○
Clontarf Pool	Estuarine	G	○
Forty Baskets Pool	Estuarine	G	○
Fairlight Beach	Estuarine	G	○
Manly Cove	Estuarine	G	○
Little Manly Cove	Estuarine	G	○
Southern Sydney – Ocean beaches			
Boat Harbour	Ocean beach	G	○
Greenhills Beach	Ocean beach	VG	○
Wanda Beach	Ocean beach	VG	○
Elouera Beach	Ocean beach	VG	○

Swimming site	Site type	Beach Suitability Grade	Change
Southern Sydney – Ocean beaches (continued)			
North Cronulla Beach	Ocean beach	VG	○
South Cronulla Beach	Ocean beach	G	↓
Shelly Beach	Ocean beach	VG	○
Oak Park	Ocean beach	VG	○
Southern Sydney – Botany Bay and lower Georges River			
Silver Beach	Estuarine	G	○
Como Baths	Estuarine	G	○
Jew Fish Bay Baths	Estuarine	P	↓
Oatley Bay Baths	Estuarine	P	↓
Carss Point Baths	Estuarine	P	↓
Sandringham Baths	Estuarine	G	○
Dolls Point Baths	Estuarine	P	↓
Ramsgate Baths	Estuarine	G	○
Monterey Baths	Estuarine	G	○
Brighton-Le-Sands Baths	Estuarine	G	○
Kyeemagh Baths	Estuarine	P	↓
Foreshores Beach	Estuarine	VP	↓
Yarra Bay	Estuarine	P	↓
Frenchmans Bay	Estuarine	P	○
Congwong Bay	Estuarine	G	○
Southern Sydney – Port Hacking			
Jibbon Beach	Estuarine	G	↓
Horderns Beach	Estuarine	G	○
GyMEA Bay Baths	Estuarine	P	○

Swimming site	Site type	Beach Suitability Grade	Change
Southern Sydney – Port Hacking (continued)			
Lilli Pilli Baths	Estuarine		
Gunnamatta Bay Baths	Estuarine		

Beach Suitability Grade					Change		
							
Very Good	Good	Fair	Poor	Very Poor	Improved	Stable	Declined

Northern Sydney (Pittwater to Manly)

Overall results

Twenty-nine of the 32 swimming sites were graded as Very Good or Good in 2021–2022. While this is a slight decline in performance on the previous year, it is still a very good result.



Percentage of sites graded as Very Good or Good

	2019–2020	2020–2021	2021–2022	Trend
Ocean beaches (20 sites)	100%	100%	100%	
Estuarine sites (10 sites)	100%	100%	90%	
Lake/lagoon sites (2 sites)	100%	0%	0%	


Beachwatch samples the ocean beaches and Narrabeen Lagoon every sixth day throughout the year, and estuarine beaches every sixth day between October and April, and monthly from May to September.

See the section on **How to read this report** on page 141 for an explanation of the graphs, tables and Beach Suitability Grades.

Best beaches

Palm Beach, Whale Beach, Avalon Beach, Mona Vale Beach, South Curl Curl Beach, Elvina Bay, The Basin and Great Mackerel Beach.

These sites had excellent water quality and were suitable for swimming almost all of the time.



Site types in Northern Sydney region

Swimming sites monitored in the Northern Sydney region include ocean beaches, estuarine areas in Pittwater and lagoon sites in Narrabeen Lagoon, with each site type having a different response to rainfall-related impacts.

Estuarine and lagoon swimming sites did not perform as well as ocean beaches due to lower levels of flushing, which increase the time needed to disperse and dilute pollution inputs, taking longer to recover from stormwater events.

As a general precaution swimming should be avoided during and for at least one day after heavy rain at ocean beaches, and for up to 3 days in estuarine and lagoon areas, or if there are signs of stormwater pollution such as discoloured water or floating debris.

Swimming is not recommended at ocean beaches located near lagoon entrances if the lagoon is open, due to the possibility of pollution from the outflow.



Beach Suitability Grades for Northern Sydney ocean beaches

Ocean beaches

All 20 ocean beaches were graded as Very Good or Good in 2021–2022.

Palm Beach, Whale Beach, Avalon Beach, Mona Vale Beach and South Curl Curl Beach were graded as Very Good. The water quality at these sites was of a very high standard and suitable for swimming almost all of the time.

Bilgola Beach, Newport Beach, Bungan Beach, Warriewood Beach, Turimetta Beach, North Narrabeen Beach, Collaroy Beach, Long Reef Beach, Dee Why, North Curl Curl Beach, Freshwater Beach, Queenscliff Beach, North Steyne Beach, South Steyne Beach and Shelly Beach were graded as Good. Water quality was frequently suitable for swimming during dry weather conditions, with elevated levels of enterococci recorded following heavy rainfall.

Bilgola Beach, Newport Beach, Bungan Beach and Dee Why Beach were downgraded to Good, due to a slight decline in microbial water quality. The decline in water quality reflects a higher proportion of samples collected at these sites during wet weather compared to the 2020–2021 assessment.

It is recommended that swimming be avoided at ocean beaches during and for up to one day following rainfall, or if there are signs of pollution such as discoloured water, flowing drains or floating debris.

Estuarine beaches

Nine of the 10 estuarine swimming sites in Pittwater were graded as Very Good or Good in 2021–2022.



Beach Suitability Grades for Northern Sydney estuarine beaches

Elvina Bay, The Basin and Great Mackerel Beach were graded as Very Good. These sites had excellent water quality and were suitable for swimming almost all of the time.

Elvina Bay was upgraded to Very Good from Good in the previous year. The microbial water quality at this site is close to the threshold between Good and Very Good and has changed between these grades over recent years. While water quality is mostly suitable for swimming during dry weather and after light to moderate rainfall, elevated bacterial levels were regularly recorded following heavy rainfall.

Barrenjoey Beach, Paradise Beach Baths, Clareville Beach, Taylors Point Baths, North Scotland Island and South Scotland Island were graded as Good. Water quality at these sites was suitable for swimming most of the time, with elevated levels of enterococci mostly recorded following rainfall. Bacterial levels were occasionally elevated at Barrenjoey Beach during dry weather conditions.

Bayview Baths was downgraded to Poor from Good, due to a decline in microbial water quality. The decline in water quality reflects a higher proportion of samples collected during wet weather conditions compared to the 2020–2021 assessment period. Elevated enterococci levels were occasionally recorded during dry weather conditions, and regularly after moderate to heavy rainfall. Water quality at this site can take longer to recover from stormwater events than at other Pittwater swimming sites due to lower levels of flushing.

Lake/lagoon swimming sites

The 2 swimming sites in Narrabeen Lagoon, Birdwood Park and Bilarong Reserve, continued to be graded as Poor in 2021–2022, as in the previous year. Water quality at these sites was mostly suitable for swimming during dry weather, with 77% and 76% of dry weather samples within the safe swimming limit for Birdwood Park and Bilarong Reserve, respectively. However, enterococci levels increased with increasing rainfall, and often exceeded the safe swimming limit after light rainfall.

Birdwood Park is located at the entrance to the lagoon and water quality at this site is influenced by wet weather events and whether the lagoon is open to the ocean. Discharge from Narrabeen Lagoon is a significant source of faecal contamination.



Beach Suitability Grades for Northern Sydney lake/lagoon swimming sites

Bilarong Reserve in Narrabeen Lagoon retains pollution inputs because it is located away from the lagoon entrance and is not well flushed by clean ocean water. A significant source of faecal contamination is stormwater runoff to the lagoon.

The amount of time the lagoon is open or closed influences water quality at Birdwood Park and Bilarong Reserve lagoon sites. During the assessment period, council undertook a major entrance clearance operation from October to December 2021, opening the entrance to the lagoon, and it has remained open as of April 2022. Prior to the entrance clearance, the lagoon entrance closed a few times between October 2019 and December 2020 and was generally closed from then on. It was mechanically opened by council on several occasions, but did not remain open for very long. Mechanical openings became more and more difficult in the lead-up to the entrance clearance as the area became more and more choked with sand. During periods of entrance closure, water quality is likely to decline as pollution inputs are not as readily dissipated or flushed.

Management



A coastal management program (CMP) outlines a long-term strategy for managing the coast, in line with the *Coastal Management Act 2016*.

The NSW Government provides guidance and funding through the Coastal and Estuary Grants Program for local councils to prepare and implement CMPs.

With funding from the NSW Government's Coastal and Estuary Grants Program, a group of councils (including Northern Beaches, Hawkesbury, Ku-ring-gai, The Hills, Central Coast and Hornsby) have completed the Hawkesbury–Nepean River System CMP Stage 1 Scoping Study. The grants program has also provided funding for Stage 2 investigations, which are currently underway and on track to be completed in 2022. The consortium is currently planning Stage 3.

The development of a CMP will allow councils to identify catchment pressures in the system (including Pittwater, Broken Bay and Brisbane Water), to prioritise management initiatives to manage issues relating to coastal and estuary health. Water quality management actions such as stormwater infrastructure improvements, restoring and maintaining riparian areas and strategic land-use planning will be considered during the process.

Under the NSW Government's Coastal and Estuary Grants Program, funding has been given to Northern Beaches Council to prepare the first stage, the scoping study, of the CMP for Collaroy–Narrabeen Beach and Fishermans Beach, and Bilgola and Mona Vale (Basin) beaches. The development of a CMP will allow the council to identify coastal hazards (which could include some water quality management actions) and prioritise initiatives to manage these.

Northern Beaches Council



North Narrabeen Beach
Photo: Beachwatch/DPE


Northern Beaches Council proactively inspects and cleans out 258 stormwater quality improvement devices. Work is continuing to standardise measurement of gross pollutants and debris and improve data management across the amalgamated council. During 2020–2021, 2 new gross pollutant traps (GPTs) were constructed, one in Manly and the other in Newport.

Northern Beaches Council's Waterway Health monitoring program is in its 12th year. The program continues to assess waterways and enables council to investigate opportunities to improve the health and amenity of these natural assets.


Water sensitive urban design has been implemented across the local government area and particularly in the Warriewood land release area, where it improves water quality prior to discharge into Narrabeen Lagoon.

Sydney Water

Sydney Water has inspected, cleaned and repaired sewer mains that have a high likelihood of discharging sewage to waterways if they become blocked. When significant tree root intrusion to the public sewer from the private sewer was identified, property owners were requested to remedy the problem.



Sampling sites and Beach Suitability Grades at Sydney's Northern Beaches



Sampling sites and Beach Suitability Grades in Pittwater

Palm Beach

Beach grade:



Palm Beach is 2.3 km long, with rock baths in the southern corner. Lifeguards patrol the beach from September to April.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and often after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Whale Beach

Beach grade: VG



Whale Beach is 600 m long, with rock baths at the southern rock platform. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Avalon Beach

Beach grade: VG



Avalon Beach is 500 m long and backed by a park and picnic area. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit in response to 20 mm or more of rain.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Bilgola Beach

Beach grade:



Bilgola Beach is 500 m long, with rock baths at the southern end. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	100%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Newport Beach

Beach grade:



Newport Beach is an open, east facing beach around 1.3 km long. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and often after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	98%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Bungan Beach

Beach grade: **G**



Bungan Beach is 600 m long and backed by a steep escarpment. Lifeguards patrol the beach from late December to the end of January.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to light rain, and often after 20 mm or more of rain.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	100%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Mona Vale Beach

Beach grade: **VG**



Mona Vale Beach is 1 km long. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 20 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Warriewood Beach

Beach grade: G



Warriewood Beach is 500 m long and located below a steep bluff. The beach is patrolled during holiday periods.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including Warriewood Wastewater Treatment Plant (WWTP).


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	96%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Turimetta Beach

Beach grade:



Turimetta Beach is 350 m long and is backed by steep bluffs. This beach is not patrolled by lifeguards.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including Warriewood WWTP.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



North Narrabeen Beach

Beach grade: **G**



North Narrabeen Beach is located at the northern end of the 3.5 km long beach and is patrolled from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from Narrabeen Lagoon.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Narrabeen Lagoon (Birdwood Park)

Beach grade: P



The Birdwood Park swimming site is a sandy beach on the southern side of the entrance to Narrabeen Lagoon. The lagoon entrance has been periodically open and closed at times.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including from elsewhere within the lagoon.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after rain.

The site has been monitored since 2004.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Lagoon	Jul 2020 to Apr 2022	77%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: D



Dry and wet weather water quality



Water quality in response to rainfall



Bilarong Reserve

Beach grade: P



Bilarong Reserve is located on the northern shoreline of Narrabeen Lagoon.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including from elsewhere within the lagoon.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to no rain, and often after rain.


See 'How to read this report' for key to map. The site has been monitored since 2014.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Lagoon	Jun 2020 to Apr 2022	76%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality




Water quality in response to rainfall



Collaroy Beach

Beach grade:



Collaroy Beach is backed by a park and picnic area. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain and often after 5 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Long Reef Beach

Beach grade:



Long Reef Beach is located near the entrance of Dee Why Lagoon. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with potential faecal contamination from discharge from Dee Why Lagoon.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and often after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: A



Dry and wet weather water quality




Water quality in response to rainfall



Dee Why Beach

Beach grade: **G**




Dee Why Beach is located at the southern end of the stretch of beach and is patrolled by lifeguards from late August to May.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to 10 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Declined 


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



North Curl Curl Beach

Beach grade: **G**



North Curl Curl Beach is located near the entrance to Curl Curl Lagoon. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from Curl Curl Lagoon.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and frequently after 20 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	96%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



South Curl Curl Beach

Beach grade: **VG**



South Curl Curl Beach is at the southern end of Curl Curl Beach and is patrolled by lifeguards from September to April.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to 10 mm or more of rain, and regularly after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Freshwater Beach

Beach grade: G



Freshwater Beach is approximately 350 m long and is patrolled by lifeguards from late August to May.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and frequently after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	96%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Queenscliff Beach

Beach grade: **G**



Queenscliff Beach is located at the northern end of Manly Beach. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination, including discharge from Manly Lagoon.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and regularly after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



North Steyne Beach

Beach grade: G



North Steyne Beach is the middle section of Manly Beach. Lifeguards patrol the beach from September to April.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater and discharge from Manly Lagoon.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and frequently after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



South Steyne Beach

Beach grade: **G**



South Steyne Beach is at the southern end of Manly Beach. Lifeguards patrol the beach year round.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after 10 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jun 2020 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Shelly Beach

Beach grade:



Shelly Beach is backed by a picnic area and reserve and is not patrolled by lifeguards.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time, but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	92%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Barrenjoey Beach

Beach grade: **G**



Barrenjoey Beach is approximately 1.5 km long and located on the north-eastern foreshore of Pittwater.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 20 mm or more.

The site has been monitored since 1996.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	91%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Paradise Beach Baths

Beach grade:



Paradise Beach Baths is a 30 by 20 m netted swimming enclosure on the eastern foreshore of Pittwater.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to light rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1996.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Clareville Beach

Beach grade:



Clareville Beach is a narrow 250 m long beach located on the eastern foreshore of Pittwater.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1995.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Taylor's Point Baths

Beach grade: **G**



Taylor's Point Baths is a 15 by 20 m netted swimming enclosure on the eastern foreshore of Pittwater.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and frequently after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 2010.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	100%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Bayview Baths

Beach grade: P



Bayview Baths is a 20 by 40 m swimming enclosure on the southern foreshore of Pittwater.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including stormwater and sewage overflows.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1995.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	89%	100	Declined ↓


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Elvina Bay

Beach grade:



Elvina Bay is located on the south-western foreshore of Pittwater. The swimming area is not netted.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels generally increased with increasing rainfall, often exceeding the safe swimming limit after 10 mm or more of rain, and regularly after 20 mm or more.

The site has been monitored since 1995.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Apr 2019 to Apr 2022	100%	100	Improved


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality




Water quality in response to rainfall



North Scotland Island

Beach grade: **G**



The North Scotland Island swimming site is a 15 by 50 m netted enclosure located on the north side of Scotland Island in Pittwater.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including onsite systems.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and often after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1995.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Mar 2019 to Apr 2022	100%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



South Scotland Island

Beach grade: **G**



The South Scotland Island swimming site is located at Carols Wharf on the southern side of Scotland Island. The location is not netted and is backed by a reserve.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including onsite systems.


Enterococci levels generally increased with increasing rainfall, often exceeding the safe swimming limit after 10 mm or more of rain, and regularly after 20 mm or more.

The site has been monitored since 1996.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Apr 2019 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



The Basin

Beach grade:



The Basin is a 500 m sandy beach on the western side of Pittwater, backed by Ku-ring-gai Chase National Park.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time with few potential sources of significant faecal contamination.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to 10 mm or more of rain, and often after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1999.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Mar 2019 to Apr 2022	98%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Great Mackerel Beach

Beach grade:



Great Mackerel Beach is a 500 m long sandy beach on the north-western side of Pittwater.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time with few potential sources of significant faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to 10 mm or more of rain, and often after 20 mm or more.

The site has been monitored since 1999.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Apr 2019 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A




Dry and wet weather water quality



Water quality in response to rainfall



Central Sydney (Bondi to Little Bay & Sydney Harbour)



Overall results

Thirty-one of the 37 swimming sites were graded as Very Good or Good in 2021–2022, a slight decline in performance from the previous year.


Percentage of sites graded as Very Good or Good

	2019–2020	2020–2021	2021–2022	Trend
Ocean beaches (10 sites)	90%	100%	90%	
Estuarine sites (26 sites)	81%	85%	81%	
Ocean baths (1 sites)	100%	100%	100%	

Beachwatch samples the ocean beaches every sixth day throughout the year, and the estuarine beaches every sixth day between October and April, and monthly from May to September.

See the section on **How to read this report** on page 141 for an explanation of the graphs, tables and Beach Suitability Grades.

Best beaches
Clovelly Beach, South Maroubra Beach and Nielsen Park.
These sites had excellent water quality and were suitable for swimming almost all of the time.



Site types in Central Sydney region

Swimming sites monitored in the Central Sydney region include ocean beaches, an ocean baths, and estuarine areas in Sydney Harbour and lower Parramatta River, with each site type having a different response to rainfall-related impacts.

Estuarine swimming sites did not perform as well as ocean beaches due to lower levels of flushing, which increase the time needed to disperse and dilute pollution inputs, taking longer to recover from stormwater events.

As a general precaution swimming should be avoided during and for at least one day after heavy rain at ocean beaches, and for up to 3 days at harbour beaches, or if there are signs of stormwater pollution such as discoloured water or floating debris.

Ocean beaches

Nine of the 10 ocean beaches were graded as Very Good or Good in 2021–2022.



Beach Suitability Grades for Central Sydney ocean beaches


Clovelly Beach and South Maroubra Beach were graded as Very Good. Water quality was suitable for swimming almost all of the time at these beaches.

Bondi Beach, Tamarama Beach, Bronte Beach, Gordons Bay, Maroubra Beach, Malabar Beach and Little Bay Beach were graded as Good. Gordons Bay and Maroubra Beach were downgraded to Good, due to a decline in microbial water quality. These sites were frequently suitable for swimming during dry weather conditions but recorded elevated enterococci levels following rainfall. Elevated enterococci levels were occasionally recorded at Malabar Beach and Little Bay Beach in dry weather conditions.

Coogee Beach was downgraded to Poor from Good in the previous year. Microbial water quality at this site has declined over the last 2 years and crossed the threshold from Good to Poor. Elevated enterococci levels were occasionally recorded during dry weather conditions and increased following rainfall. The decline in water quality reflects a higher proportion of samples collected during wet weather compared to the 2020–2021 assessment.

It is recommended that swimming be avoided at ocean beaches during and for up to one day following rainfall, or if there are signs of pollution such as discoloured water, flowing drains or floating debris.

Estuarine beaches



Beach Suitability Grades for Central Sydney estuarine beaches

Twenty-one of the 26 estuarine swimming sites in Sydney Harbour were graded as Very Good or Good in 2021–2022.

Nielsen Park continued to be graded as Very Good, as in previous years. This site has excellent water quality, with few potential sources of faecal contamination, and is suitable for swimming almost all of the time. The site is close to the open ocean allowing any pollution inputs to be quickly diluted and dispersed.

Twenty of the estuarine swimming sites in Sydney Harbour were graded as Good: Camp Cove, Watsons Bay, Parsley Bay, Murray Rose Pool, Dawn Fraser Pool, Chiswick Baths, Cabarita Beach, Woolwich Baths, Woodford Bay, Greenwich Baths, Hayes St Beach, Clifton Gardens, Balmoral Baths, Edwards Beach, Chinamans Beach, Clontarf Pool, Forty Baskets Pool, Fairlight Beach, Manly Cove and Little Manly Cove. Camp Cove was downgraded to Good, due to a slight decline in microbial water quality. These sites had mostly good water quality, although enterococci levels increased following rainfall.

Gurney Crescent Baths continued to be graded Fair, as in previous years. Microbial water quality at this site is frequently suitable for swimming during dry weather conditions, with 93% of dry weather samples within the safe swimming limit. However, enterococci levels increase following rainfall. This site has more significant sources of microbial contamination including upstream sources in Middle Harbour.

Four estuarine swimming sites in Sydney Harbour were graded as Poor in 2021–2022: Rose Bay Beach, Tambourine Bay, Northbridge Baths and Davidson Reserve. Water quality at these swimming sites was mostly suitable for swimming during dry weather conditions, with elevated bacterial levels recorded following rainfall.

While Tambourine Bay in Lane Cove River, and Davidson Reserve in Middle Harbour continued to be graded as Poor from the previous year, Rose Bay Beach in Port Jackson and Northbridge Baths were downgraded to Poor from Good and Fair grades, respectively, due to a decline in microbial water quality. The decline in microbial water quality reflects a higher proportion of samples collected during wet weather at these sites compared to the 2020–2021 assessment. These sites are more susceptible to wet weather impacts and have several significant sources of faecal contamination including upstream sources, stormwater and sewage overflows. During late February, March and early April 2022, extreme wet weather led the sewage overflow discharge capture capacity of Sydney Water’s Northside Storage Tunnel to be

exceeded. As a result, discharges occurred to Middle Harbour and Lane Cove River. Since 2000, when the tunnel was included in the system, it has performed flawlessly preventing large volumes of sewage overflow from entering these waterways, however it has limits, and these were exceeded due to the unprecedented combination of widespread intense rainfall and prolonged duration. Sewage overflows are hydraulic relief points included on wastewater systems to prevent sewage backing up and surcharging to homes.

Estuarine sites are not as well flushed as ocean beaches, and so can take longer to recover from stormwater events. As a precaution, swimming should be avoided at Sydney Harbour swimming sites during and for up to 3 days following rainfall or if there are signs of pollution such as discoloured water, flowing stormwater drains or floating debris.



Beach Suitability Grades for Central Sydney ocean baths

Ocean baths

South Maroubra Rockpool was graded as Good in 2021–2022, similar to previous years. Water quality is frequently suitable for swimming during dry weather conditions, with 94% of dry weather samples within the safe swimming limit. Enterococci levels increased with increasing rainfall. Swimming should be avoided during and for up to one day following rainfall, or if there are any signs of pollution such as the stormwater drain discharging to the site, discoloured water or floating debris.

Management

Ocean beaches

In 2019, the NSW Government committed \$2.5 million for the diversion of stormwater from Coogee Beach to improve water quality and the marine environment. The intention of the Government is that all payments, including those for structural works, be subsequent to negotiation and agreement to the project design by the Coogee Beach Working Group.



With funding from the NSW Government's Coastal and Estuary Grants Program, Woollahra, Waverley and Randwick City councils have prepared the first stage of the Eastern Sydney Beaches CMP, the scoping study. The development of a CMP will allow the councils to identify coastal hazards (which could include some water quality management actions) and prioritise initiatives to manage these.

Waverley Council

A coastal management program (CMP) outlines a long-term strategy for managing the coast, in line with the *Coastal Management Act 2016*.

The NSW Government provides guidance and funding through the Coastal and Estuary Grants Program for local councils to prepare and implement CMPs.

Council is working with Sydney Water to progress the Refresh Vaucluse and Diamond Bay project, which will connect the suburbs of Watsons Bay and Vaucluse and parts of Dover Heights to the Bondi Water Resource Recovery Facility (WRRF) to divert wastewater flows from the last 3 remaining ocean outfalls in Sydney. The pipeline will be located behind properties along Kimberley Street, near the existing boardwalk. Construction is set to commence in early 2023.


GPTs have been installed in the Bondi Beach and Bronte Beach catchments, and the North Bondi GPT is currently undergoing an upgrade to improve its effectiveness.

The Bronte Stormwater Harvesting Scheme collects and treats stormwater, which is then re-used for toilets, park irrigation and ocean pool cleaning. The scheme saves over 16 million litres of water each year and reduces the volume of stormwater discharged to Bronte Beach.

The Bondi Stormwater Harvesting Scheme commenced in 2012 and supplies approximately 50 million litres of treated stormwater for park irrigation and toilets in Bondi Pavilion and South Bondi. The restored Bondi Pavilion will be connected to the scheme, which will potentially increase the use of the recycled water resources in Bondi Beach. An underground filtration system is installed that treats excess stormwater runoff from Campbell Parade, resulting in cleaner water at Bondi Beach.

The Tamarama Stormwater Harvesting Scheme commenced operation in December 2015 supplying treated stormwater for park irrigation and toilets in Tamarama Park. The scheme supplies approximately 14 million litres of water each year and reduces the volume of stormwater discharged to Tamarama Beach. As part of the scheme, a large capacity underground sediment basin has also been installed to prevent sediment and other pollutants entering the ocean at Tamarama Beach.

Randwick City Council



Gordons Bay

Photo: Beachwatch/DPE

Randwick City Council operates and maintains 15 stormwater harvesting treatment systems with UV filtration across the local government area. These systems treat stormwater by removing suspended solids, bacteria and other organic and inorganic materials before it is used for irrigation in surrounding landscaped and garden areas, reducing stormwater discharging to the beaches.

In 2021, Randwick City Council completed construction of 2 stormwater harvesting systems; one located at Maroubra foreshore to improve the water quality at Maroubra Beach and South Maroubra Rockpool, and one at Purcell Park in Matraville.

Randwick City Council maintains 36 GPTs on stormwater lines leading to the local bays, which are all cleaned regularly. From February 2021 to February 2022, approximately 372 tonnes of material was removed from these GPTs. There is also a systematic cleaning program for all drainage pits including a regular street sweeping program that assists with reducing stormwater pollution to the local bays. Randwick City Council continues to conduct litter education campaigns throughout the local government area to educate residents on the proper disposal of waste. This program aims to reduce the amount of litter disposed on beaches and entering the ocean.

Randwick City Council has a strategic program and reactive process to monitor and assess the condition of the stormwater pipes in the local area using CCTV.

The Coogee Beach Stormwater Quality Working Group was established in 2017 to improve water quality at Coogee Beach. The group consists of representatives from the community, Sydney Water, Beachwatch, council staff and the State Member for Coogee. In August 2017, the working group put recommendations to council to investigate the feasibility of diverting more stormwater from the beach, and the development of a community education and marketing campaign to better inform the local community about stormwater and its impact on local beaches. In May 2022, with funding from the NSW Department of Planning and Environment (DPE), a consultant was appointed and will meet with the Coogee Beach Stormwater Quality Working Group to investigate and design (including water harvesting assessment) a solution for stormwater diversion at Coogee Beach.

Council officers undertake their routine inspections and regulatory duties to ensure stormwater pollution is investigated and mitigated to reduce impacts to the water quality of local recreational waterways.

Sydney Water

Sydney Water investigated wet weather sewage overflows in the Coogee Beach catchment and found that silt was accumulating within the Coogee Diversion Sewer (CDS) due to the very low slope of this sewer. As a result, Sydney Water is undertaking more frequent de-silting of the CDS and the grit pits at the northern end of the beach. This work will increase the capacity of the CDS and reduce the occurrence of overflows.

Sydney Water has inspected, cleaned and repaired sewer mains that have a high likelihood of discharging sewage to waterways if they become blocked. When significant tree root intrusion to the public sewer from the private sewer was identified, property owners were requested to remedy the problem.

Sydney Harbour



Sampling at Rose Bay
Photo: Beachwatch/DPE

In 2019, the NSW Government committed \$150,000 to address the poor recreational water quality at Rose Bay Beach. DPE, in collaboration with the University of Technology Sydney, conducted a catchment investigation applying enterococci and genetic marker assay methods for a detailed audit of sources of contamination. The project identified that microbial pollution in Rose Bay is largely from human sewage, which contaminates stormwater particularly after rain. In September 2020, the findings of the investigation were released in *Towards Safer Swimming: Rose Bay*, and are being used to focus remediation efforts in the catchment and help design and implement management strategies to resolve water quality issues at the beach.

Under the leadership of Greater Sydney Local Land Services, a consortium of state agencies, Sydney Water and 17 Sydney Harbour councils prepared the Greater Sydney Harbour CMP Stage 1 Scoping Study and have also completed the Greater Sydney Harbour Estuary Processes Study, which contributes to Stage 2 of the CMP process and will inform Stage 3. Both studies were supported with funding from the NSW Government's Coastal and Estuary Grants Program.

The Sydney Coastal Councils Group (SCCG) in partnership with the Parramatta River Catchment Group (PRCG), together with a working group made up of representatives from Sydney Water and DPE, are leading the development of the remainder of the CMP. Stage 2 investigations are currently underway and expected to be completed by the end of 2022. The program will identify catchment pressures and prioritise management initiatives for issues relating to coastal and estuary health. Water quality management actions such

as stormwater infrastructure improvements, restoring and maintaining riparian areas, and strategic land-use planning will be considered during the process.

The Lane Cove River Coastal Zone Management Plan (CZMP) is implemented by local councils including Lane Cove, Hunters Hill, Ryde and Willoughby City.

Parramatta River Estuary CZMP is being implemented by several councils including Cumberland City, (formerly Auburn), City of Canada Bay, Hunters Hill, Parramatta City, Inner West (formerly Ashfield and Leichhardt), City of Ryde and Strathfield. With funding from the NSW Government's Coastal and Estuary Grants Program, many of the actions from the CZMP have been completed, including the installation of a GPT and bush regeneration to improve water quality in Tarban Creek in the Hunters Hill local government area. Further sediment and weed removal work is being undertaken at Tarban Creek to remediate the coastal saltmarsh and mangroves.

Northern Beaches Council



Manly Cove

Photo: Beachwatch/DPE

Northern Beaches Council proactively inspects and cleans out 258 stormwater quality improvement devices in the local government area. Work is continuing to standardise measurement of gross pollutants and debris and improve data management across the amalgamated council.

Northern Beaches Council maintains a constructed raingarden within East Esplanade Reserve. The raingarden eases localised flooding and removes nutrient loads from stormwater before it enters Manly Cove. In 2020–2021 council undertook significant landscaping of East Esplanade Reserve, which dramatically reduced the amount of runoff and erosion from the reserve entering Manly Cove.

Mosman Council

Mosman Council's Botanic Road Stormwater Re-use Scheme is an underground storage system that captures stormwater and provides UV disinfection, after which it is pumped to Balmoral Oval and Balmoral Reserve for irrigation. This reduces the amount of stormwater entering our waterways.

There are over 30 stormwater quality improvement devices throughout Mosman, installed at Balmoral Beach, Clifton Gardens, Edwards Beach and Chinamans Beach, to capture sediment and floating debris. Council has installed stormwater quality improvement devices at more than 75% of stormwater outlet points. These devices capture sediment

and floating debris from stormwater before it enters the waterways.

Council continues to maintain natural riparian areas to protect vegetation and improve water quality in Sydney Harbour. Recent improvement works have been completed at Clifton Gardens with sediment removal from the creek lines and ponds installed to reduce flooding and stormwater runoff into Sydney Harbour. The works also included the revegetation of the creek banks with native sedges and other planting work using local native species.



Clifton Gardens

Photo: Beachwatch/DPE

Council undertakes regular cleaning activities at all council beaches, parks, reserves and ovals. Litter is removed by clearing contractors from the foreshore, and by using mechanical beach rakes and hand raking. Council's street sweeping program regularly cleans roads, gutters and footpaths using mechanical street sweepers and manual sweeping. These activities aim to prevent litter and waste reaching council's beaches and waterways.

Mosman Council continues to implement its HarbourCare volunteer program. The program helps residents to be actively involved in collecting rubbish along the Mosman foreshore. Data on the rubbish collected is collated and reported back to council. Mosman Council then uses this data to help in the development of education programs.

Council regularly runs campaigns and events promoting litter reduction, specially targeting cigarette butts, fishing tackle and outdoor party litter. In September 2022, the first Seaside Scavenge event run by Take 3 was held at Balmoral Reserve. The litter was sorted, counted and logged on the Tangaroa Blue Foundation database. Six new Department of Primary Industries tackle bins have been reintroduced to Clifton Gardens and Balmoral wharf and Spit West, which are regularly used.

Over 26 groups participated in the 2022 Clean Up Australia Day, with some of the groups targeting litter collection at Chinamans Beach, Balmoral Beach, Clifton Gardens and Rawson Oval.

Willoughby City Council

Willoughby City Council has signage at Northbridge Baths to advise the community not to swim during and for up to 48 hours after rainfall due to potential stormwater pollution.


Council has a water plan with innovative stormwater cleansing actions including targeted pollution reduction filtration pits, stormwater biofiltration interventions, ponded wetlands and GPTs. Council has included funding for

maintenance of these stormwater cleansing assets to ensure consistent reliability in their long-term financial plan.

Willoughby City Council has ongoing water quality monitoring in all creeks within the local government area. It has funding for stormwater projects including stormwater harvesting, stormwater quality improvement devices, and to complete a detailed design for a future biofiltration ponded wetland and stormwater harvesting project at Bicentennial Oval. This will complement council's existing stormwater harvesting plant at Artarmon reserve.

North Sydney Council

In April 2022 North Sydney Council installed 2 new GPTs to help improve the water quality in the harbour and prevent sediments and litter entering the waterways. These included one in Brennan Park and another in Newlands Lane, Wollstonecraft.



Hayes St Beach

Photo: Beachwatch/DPE

Council undertakes regular catchment water quality monitoring. It also supports beach, foreshore and water clean-ups, including through HarbourCare volunteers, who are concerned about pollution in Sydney Harbour and its effect on marine and bird life.

North Sydney Council's Stormwater Re-use Project continues to harvest, treat and re-use stormwater for the irrigation of sports fields and recreational parks, including St Leonards Park, Cammeray Park, Forsyth Park, Primrose Park and Tunks Park. This saves millions of litres of potable water, improves the quality and reduces the amount of stormwater entering the waterways.

Lane Cove Council

Lane Cove Council maintains 14 GPTs in the catchment to reduce the impact of stormwater to the waterways. In 2021–2022 more than 80 tonnes of material was prevented from entering the Lane Cove River. Over 80% of this material was recycled.

Lane Cove Council is an active participant in the Greater Sydney Harbour CMP – Stage 2 Planning Grant.

With funding from the NSW Government's Coastal and Estuary Grants Program, Lane Cove Council completed Stage 2 of the stormwater improvement works at Lane Cove Bushland Park in 2020, an action from the Lane Cove CZMP. The work has improved 3 drainage lines into Gore Creek, stabilised the creek bed, and protected endangered ecological communities in the vicinity.

Hunters Hill Council

Hunters Hill Council maintains 24 stormwater quality improvement devices in the local government area. Council has assessed the condition of 20% of stormwater assets as part of its Asset Management Plan. In addition, council is ensuring stormwater assets are routinely inspected and maintained into the future.

City of Canada Bay Council

The City of Canada Bay maintains over 27 stormwater quality improvement devices that prevent over 150 tonnes of pollutants (sediments, leaves and litter) from reaching the Parramatta River each year. Stormwater harvesting, rainwater re-use and raingardens have been constructed in the Drummoyne Oval precinct to reduce stormwater and pollutant loads reaching Five Dock Bay.



Chiswick Baths

Photo: Beachwatch/DPE

The *Our Water for Our Community* stormwater recycling scheme at Cintra Park, completed in October 2015, harvests and re-uses stormwater for the irrigation of 2 public golf courses and 15 sporting fields. This reduces the City of Canada Bay's reliance on potable water by 180 million litres each year, and improves the quality and reduces the quantity of runoff into Canada Bay itself.

As part of the Parramatta River Catchment Group (PRCG) mission to make the Parramatta River swimmable again, the City of Canada Bay is in the planning stages of activating 2 new sites to provide greater access to the river. At Bayview Park, Concord Council is planning to reintroduce a netted swimming enclosure adjacent to the natural sandy beach. Replacement of the seawall with a new environmentally friendly design at McIlwaine Park, Rhodes will provide the opportunity for splash contact in newly constructed rock pools and provision of launch facilities for water activities such as kayaking, rowing and stand-up paddle boarding.

Woollahra Municipal Council

Woollahra Municipal Council undertakes a range of projects to improve water quality at its harbour beaches. Council is continually upgrading stormwater infrastructure, installing and maintaining GPTs and raingardens to remove contaminants from stormwater. Street sweeping, beach cleaning, riparian vegetation and terrestrial bushland regeneration activities continue to contribute to improved stormwater quality at Woollahra's beaches.

In 2022 council installed 2 new GPTs, one in Woollahra, treating the Double Bay catchment, and the other in Vaucluse, treating the Parsley Bay catchment. A number of

pit litter baskets were replaced in the Double Bay catchment, to improve litter capture.

Council continued upgrading raingardens throughout the municipality, replacing filtration media and plants to ensure that pollutant capture is maximised. The raingardens capture and treat first-flush stormwater runoff.



Camp Cove

Photo: Beachwatch/DPE

Woollahra Municipal Council is a member of the Rose Bay Beach Working Party that was established in 2017 to address water quality issues at Rose Bay Beach. Other members of the working party include representatives from DPE, Transport for NSW (Maritime), Sydney Water and the Member for Vaucluse. The Working Party has identified the sources of pollution at Rose Bay Beach and continues to work in partnership to address them. This has included council and Sydney Water working together to identify and resolve a number of sewer chokes and leaks.

Also in the Rose Bay catchment, council has proceeded to detailed design on a project to improve the stormwater outlet at Caledonian Road. The GPT on that outlet is functioning well, however there remains an issue of scour and ponding. The project under design will see the installation of scour protection and redirection of low flows of stormwater.

Council undertakes a range of programs to educate the community about ways they can improve water quality, from picking up dog waste to undertaking water sensitive urban design. In 2021–2022 council held a number of 'rockpool rambles' with a marine ecologist, highlighting the importance of keeping Sydney Harbour as a habitat and educating the community about how they can prevent stormwater pollution. Council also stencilled a number of drains in the Rose Bay commercial centre, reminding people that the drains flow to Rose Bay Beach.

Council continues to support community groups, such as Splash without the Trash, to undertake beach clean-ups and also continues to have strong interest in the HarbourCare program, which supports volunteers to undertake clean-ups of beachside areas.

Council is participating in the Greater Sydney Harbour CMP, which has a strong focus on improving water quality harbour-wide.

Inner West Council

Inner West Council owns and maintains several stormwater quality improvement features including raingardens, constructed wetlands and GPTs to filter and clean

stormwater runoff in the catchments before discharging to Parramatta River, Sydney Harbour and the Cooks River.

Council continues to develop green-blue grid planning across its local government area to create further areas of green space and water sensitive design as well as increase local access to council's coastal areas. To support this work, council continues to work as part of the Cooks River Alliance and Parramatta River Catchment Group to support increased water quality endeavours, and is contributing to the current review and update of the CMP for these 2 river systems and Sydney Harbour.



Dawn Fraser Pool

Photo: Beachwatch/DPE

Inner West Council currently has 26 GPTs across its local government area and has been working with external contractors to restore these units to full working order and avoid litter bypassing the units and entering the harbour. Council is reviewing the design and viability of 4 new GPTs in locations across the Parramatta River, for construction in future years. In many of council's waterfront parks, these units are also being considered as the first step for future stormwater harvesting schemes for field irrigation. To support and incentivise the community take-up of water re-use, Inner West Council provides a rainwater tank rebate, and workshops for rainwater harvesting and water sensitive urban design on your property. The workshops include information on reducing potable water use and rainwater, stormwater and greywater re-use.

Parramatta River Catchment Group

The Parramatta River Catchment Group (PRCG) is an alliance of councils, government agencies and community groups working together to transform the Parramatta River and its tributaries into living waterways. The group has produced a 10-step masterplan (10 recommendations) for improving the suitability of the river for swimming and its ecosystem health. By addressing water quality impacts such as stormwater run-off and sewage overflows in a coordinated way across local government areas, it is expected that more sites, including riverside beaches, will meet requirements to enable safe swimming along the river.

Sydney Water

Sydney Water is the lead coordinating agency driving the delivery of the PRCG's Parramatta River Masterplan. Sydney Water is implementing the masterplan's Step 6 to improve overflows, by undertaking a substantial sewer inspection and improvement program across the catchments of upper Parramatta River, Duck Creek and the Lane Cove River. This will reduce the inflow of stormwater to the wastewater

system and should lessen the impact of sewage overflows to these waterways.

Sydney Water is progressing a project to reduce the occurrence of wastewater overflows discharging to Rose Bay, Port Jackson. This work is an action from the Rose Bay Beach Working Party and will assist with improving the water quality at the beach.



Cabarita Beach
Photo: Beachwatch/DPE


Sydney Water has inspected, cleaned and repaired sewer mains on the northern and southern sides of Port Jackson that have a high likelihood of discharging sewage to waterways if they become blocked. Where significant tree root intrusion to the public sewer from the private sewer was identified, property owners were requested to remedy the problem.

Environment Protection Authority

The NSW Environment Protection Authority (EPA) is the primary environmental regulator for NSW. The EPA is committed to improved water quality outcomes across NSW and works closely with the PRCG to implement the masterplan for the Parramatta River. The EPA undertakes a range of regulatory functions within the Parramatta River catchment including the regulation of industrial premises, contaminated sites and significant infrastructure, and through membership of the PRCG provides advice to councils on the activation of swim sites along the Parramatta River. The EPA also participates in the PRCG's annual Get the Site Right campaign, a month long blitz of inspections of construction sites across the Parramatta River catchment, designed to reduce sediment entering waterways as pollution from building sites. This is particularly relevant for the estuarine beaches in the Sydney region, which can be heavily impacted by poor water quality following rainfall events. In this year's campaign, the EPA inspected 21 sites it regulates and accompanied local Sydney councils inspecting an additional 51 building sites.



Sampling sites and Beach Suitability Grades at Sydney's central beaches



Sampling sites and Beach Suitability Grades in Sydney Harbour

Bondi Beach

Beach grade: **G**



Bondi Beach is 800 m long and backed by a promenade, carpark and parklands, and lifeguards patrol the beach year round.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Tamarama Beach

Beach grade: G



Tamarama Beach is approximately 80 m long and lifeguards patrol the beach from late September to April.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often in response to 10 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	96%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Bronte Beach

Beach grade: **G**



Bronte Beach is 250 m long and backed by a large park and picnic area. Lifeguards patrol the beach from September to May.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	94%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Clovelly Beach

Beach grade:



Clovelly Beach is at the end of a long and narrow bay and is protected from ocean swells, and is patrolled from late September to April.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 10 mm or more of rain, and often after 20 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	96%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Gordons Bay

Beach grade:



Gordons Bay is long and narrow with a small beach located at the end of the bay and is not patrolled by lifeguards.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 2013.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	100%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Coogee Beach

Beach grade: P



Coogee Beach is 400 m long and is backed by a promenade and parklands and is patrolled by lifeguards all year round.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	87%	100	Declined ↓


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Maroubra Beach

Beach grade: **G**

Maroubra Beach is 1 km long and lifeguards patrol the beach all year round.



The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and often after 20 mm or more.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	93%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



South Maroubra Beach

Beach grade: **VG**



South Maroubra Beach is located at the southern end of Maroubra Beach and lifeguards patrol the beach all year round.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.


Enterococci levels generally increased with increasing rainfall, often exceeding the safe swimming limit after 5 mm or more of rain.

The site has been monitored since 2012.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



South Maroubra Rockpool

Beach grade: **G**



South Maroubra Rockpool is located at the southern end of Maroubra Beach and is not patrolled. During very low tides, the rockpool may be empty.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 2012.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean baths	Aug 2020 to Apr 2022	94%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Malabar Beach

Beach grade: G



Malabar Beach is 150 m long and located at the end of a long, narrow bay and is not patrolled by lifeguards.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	93%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Little Bay Beach

Beach grade: G



Little Bay Beach is a small, crescent-shaped beach bounded by rocky headlands to the north and south and is not patrolled.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site was monitored from 1989 until 1995, and since 2006.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	89%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Camp Cove

Beach grade:



The Camp Cove swimming area is not netted and is backed by a narrow stretch of beach. Lifeguards patrol this swimming site during the summer period.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit in response to 20 mm or more of rain.


See 'How to read this report' for key to map. The site was monitored since 2015.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	94%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Watsons Bay

Beach grade:



The swimming site is a 20 by 40 m enclosed tidal swimming area with a narrow sandy beach and is backed by parklands with picnic facilities.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Parsley Bay

Beach grade: **G**



The swimming site is a netted swimming area backed by a sandy beach and reserve with picnic facilities and a playground.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Nielsen Park

Beach grade: **VG**



Nielsen Park swimming area is approximately 150 m long and is netted from October to April. It is backed by a sandy beach and Sydney Harbour National Park.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with several potential sources of minor faecal contamination.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after rainfall.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	97%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Rose Bay Beach

Beach grade: P



Rose Bay Beach is approximately 500 m long and the swimming area is not netted.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including stormwater and sewer chokes.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	82%	100	Declined ↓


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality




Water quality in response to rainfall



Murray Rose Pool

Beach grade:



Murray Rose Pool (formerly Redleaf Pool) is a netted swimming enclosure in Double Bay, at the end of Seven Shillings Beach.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	97%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Dawn Fraser Pool

Beach grade:



Dawn Fraser Pool is an enclosed swimming area located in the Parramatta River and is open between October and April each year.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream sources in the Parramatta River.


Enterococci levels increased with increasing rainfall, often exceeding the safe swimming limit after 5 mm or more of rain, and frequently after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	94%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Chiswick Baths

Beach grade:



Chiswick Baths is a netted swimming enclosure in Five Dock Bay and is backed by a narrow sandy beach and a park.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream sources in the Parramatta River.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and regularly after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1999.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	92%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Cabarita Beach

Beach grade:



Cabarita Beach is a 120 m long sandy beach and is backed by parklands.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with potential faecal contamination from sewage overflows and upstream sources in the Parramatta River.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to light rain, and often after 10 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1996.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	93%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Woolwich Baths

Beach grade:



Woolwich Baths is a 20 by 30 m netted swimming area in the lower Lane Cove River with a narrow sandy beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with potential faecal contamination from stormwater and upstream river sources.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	88%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Tambourine Bay



Beach grade:



Tambourine Bay is in the lower Lane Cove River. The swimming enclosure has been removed and access to the water is limited.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with potential faecal contamination including stormwater and upstream river sources.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	90%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Woodford Bay

Beach grade: **G**



This site is a 20 by 25 m swimming enclosure on the western side of Woodford Bay in the lower Lane Cove River.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from upstream river sources.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	90%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Greenwich Baths

Beach grade: **G**



Greenwich Baths is a 40 m long netted swimming area backed by a sandy beach and is open during the swimming season.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from upstream river sources.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and frequently after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	86%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Hayes Street Beach

Beach grade: **G**

Hayes Street Beach is approximately 50 m long and is located adjacent to the Hayes Street Ferry Wharf in Neutral Bay and is not netted.



The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

The site has been monitored since 1994.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Clifton Gardens

Beach grade: **G**



Clifton Gardens is a large netted swimming area at the western end of a 250 m long beach in Chowder Bay and is backed by Sydney Harbour National Park and a park.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Jul 2019 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Balmoral Baths

Beach grade: G



Balmoral Baths is a netted swimming area at the eastern end of Balmoral Beach and is backed by a park.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Jul 2019 to Apr 2022	98%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Edwards Beach

Beach grade: **G**



Edwards Beach is a popular swimming area backed by a walking track, park and café facilities.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels increased with increasing rainfall, often exceeding the safe swimming limit after 10 mm or more of rain.

The site has been monitored since 1994.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	97%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Chinamans Beach

Beach grade: **G**



Chinamans Beach is approximately 250 m long and is a popular swimming area in Middle Harbour. It is backed by Rosherville Reserve.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream sources in Middle Harbour.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1998.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	95%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Northbridge Baths

Beach grade: P



Northbridge Baths is a 30 by 65 m enclosed swimming area in Sailors Bay, Middle Harbour and is open year round.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including stormwater and upstream sources in Middle Harbour.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to no rain, and often after light rain.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	92%	100	Declined ↓


Sanitary inspection: High




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Davidson Reserve



Beach grade:



Davidson Reserve is a 25 m long swimming area situated in Middle Harbour and is backed by Garigal National Park and picnic area.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including sewage overflows and upstream sources in Middle Harbour.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain and regularly after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	94%	100	Stable


Sanitary inspection: High




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Gurney Crescent Baths

Beach grade: **F**



Gurney Crescent Baths is a 20 m square netted swimming area located at Pickering Point in Middle Harbour.


The Beach Suitability Grade of Fair indicates microbial water quality is occasionally susceptible to faecal pollution, usually triggered by rainfall, with several potential sources of faecal contamination including upstream sources in Middle Harbour and stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1996.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	93%	100	Stable


Sanitary inspection: High




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Clontarf Pool

Beach grade:



Clontarf Pool is a small netted swimming area in Middle Harbour backed by a narrow sandy beach and a park.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream sources in Middle Harbour and stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	93%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Forty Baskets Pool

Beach grade: **G**



Forty Baskets Pool is a 20 by 40 m netted swimming area at the northern end of Forty Baskets Beach in North Harbour.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to 5 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	97%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Fairlight Beach

Beach grade:



Fairlight Beach is a narrow beach located in North Harbour. A 25 m pool filled with water from the harbour is adjacent to the beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1996.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	94%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Manly Cove

Beach grade:



Manly Cove is a netted swimming enclosure at the centre of the 250 m long beach, adjacent to the Manly Ferry Terminal.



The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and frequently after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	85%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Little Manly Cove

Beach grade: **G**



The 30 m square swimming enclosure is at the eastern end of the sandy beach in Little Manly Cove.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

The site has been monitored since 1994.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	91%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Southern Sydney (Sutherland beaches, lower Georges River, Botany Bay & Port Hacking)



Overall results

Nineteen of the 28 swimming sites were graded as Very Good or Good in 2021–2022. This is decline in performance from the previous year.

Percentage of sites graded as Very Good or Good

	2019-2020	2020-2021	2021-2022	Trend
Ocean beaches (8 sites)	100%	100%	100%	
Estuarine sites (20 sites)	90%	85%	55%	


Beachwatch samples the ocean beaches every sixth day throughout the year, and estuarine beaches every sixth day between October and April, and monthly from May to September.

See the section on **How to read this report** on page 141 for an explanation of the graphs, tables and Beach Suitability Grades.

Best beaches

Greenhills Beach, Wanda Beach, Elouera Beach, North Cronulla Beach, Shelly Beach and Oak Park.

These sites had excellent water quality and were suitable for swimming almost all of the time.



Site types in Southern Sydney region

Swimming sites monitored in the Southern Sydney region include ocean beaches and estuarine areas in Botany Bay, lower Georges River and Port Hacking, with each site type having a different response to rainfall-related impacts.

Estuarine swimming sites did not perform as well as ocean beaches due to lower levels of flushing, which increases the time needed to disperse and dilute pollution inputs, taking longer to recover from stormwater events.

As a general precaution swimming should be avoided during and for at least one day after heavy rain at ocean beaches, and for up to 3 days in estuarine areas, or if there are signs of stormwater pollution such as discoloured water or floating debris.

Ocean beaches

All 8 Southern Sydney ocean beaches were graded as Very Good or Good in 2021–2022.




Beach Suitability Grades for Southern Sydney ocean beaches

Greenhills Beach, Wanda Beach, Elouera Beach, North Cronulla Beach, Shelly Beach and Oak Park were graded as Very Good. Water quality at these sites has been consistently excellent for many years and is suitable for swimming almost all of the time.

Boat Harbour and South Cronulla Beach were graded Good. South Cronulla Beach was downgraded from Very Good in the previous year due to a slight decline in microbial water quality. Water quality at these sites is mostly suitable for swimming during dry weather conditions, but can be susceptible to pollution, with elevated enterococci levels occasionally recorded during dry weather or after light rainfall.

It is recommended that swimming be avoided at ocean beaches during and for up to one day following rainfall, or if there are signs of pollution such as discoloured water, flowing drains or floating debris.

Estuarine beaches



Beach Suitability Grades for Southern Sydney estuarine beaches

Eleven of the 20 estuarine sites were graded as Good in 2021–2022: Silver Beach, Como Baths, Sandringham Baths, Ramsgate Baths, Monterey Baths, Brighton-Le-Sands Baths, Congwong Bay, Jibbon Beach, Horderns Beach, Lilli Pilli Baths and Gunnamatta Bay Baths. Water quality at these sites was suitable for swimming most of the time, with elevated levels of enterococci mostly recorded following rainfall.

Eight estuarine swimming sites were graded as Poor in 2021–2022: Jew Fish Bay Baths, Oatley Bay Baths and Carss Point Baths in the lower Georges River, Dolls Point Baths, Kyeemagh Baths, Frenchmans Bay and Yarra Bay in Botany Bay, and Gynea Bay Baths in Port Hacking. Microbial water quality at these sites was mostly suitable for swimming during dry weather conditions, with between 77% and 95% of dry weather samples within the safe swimming limit. However, bacterial levels increased significantly with increasing rainfall.

With the exception of Frenchmans Bay and Gynea Bay Baths, these sites were all downgraded from Good in the previous year. The decline in microbial water quality at these sites reflects a higher proportion of samples collected during wet weather at these sites compared to the 2020–2021 assessment. These sites can be susceptible to wet weather impacts and have several sources of faecal contamination including upstream sources, stormwater and sewage overflows and chokes.

Foreshores Beach was downgraded to Very Poor from Poor in the previous year. Water quality at this site is significantly impacted by faecal contamination during and following rainfall, and occasionally during dry weather. The site is very susceptible to faecal contamination from the sewage overflows that periodically discharge into Mill Stream. Sydney Water have placed permanent signage to advise the public to avoid swimming 3 days after rainfall due to the risk from sewage overflows which may impact water quality at this site.

Swimming should be avoided during and for up to 3 days following light rainfall at all estuarine swimming sites, or if there are signs of pollution such as discoloured water, odours or floating debris.

Management

Ocean beaches



Under the NSW Government's Coastal and Estuary Grants Program, funding has been given to Sutherland Shire Council to prepare the Bate Bay CMP. The program will identify coastal hazards (which could include some water quality management actions) and prioritise initiatives to manage these.

Sutherland Shire Council

Sutherland Shire Council continues to see the ongoing benefits from the Cronulla-Woolooware Wastewater Reuse Scheme (CWWRS). The facility can recycle up to 4 million litres of treated wastewater each day, which is used to irrigate ovals and recreation areas across the Sutherland Shire. This includes public and private sites such as Greenhills Marang Parklands, Cronulla Golf Course, Woolooware High School, Woolooware Golf Course, Captain Cook Playing Fields and the Solander Playing Fields. The CWWRS has reduced the demand for potable water for irrigation purposes and reduced the amount of treated effluent discharged to ocean outfalls.



Greenhills

Photo: Beachwatch/DPE

In April 2022 Sutherland Shire Council adopted the Bate Bay CMP, prepared in accordance with the NSW Government's coastal management framework. Although Bate Bay water quality was found to be consistently good, the CMP recommends council continue to pursue opportunities to manage stormwater discharges to the bay through installation of stormwater quality improvement devices, community education and enforcement, and ongoing collaboration with local community groups on environmental management initiatives.

Sydney Water

To reduce the incidence of wet weather sewage overflows in beach catchments across the Cronulla Peninsula, Sydney Water has amplified sewer pipes and pumps and included storage tanks.

Sydney Water has inspected, cleaned and repaired sewer mains that have a high likelihood of discharging sewage to waterways if they become blocked. When significant tree root intrusion to the public sewer from the private sewer was identified, property owners were requested to remedy the problem.

Botany Bay and lower Georges River

A coastal management program (CMP) outlines a long-term strategy for managing the coast, in line with the *Coastal Management Act 2016*.

The NSW Government provides guidance and funding through the Coastal and Estuary Grants Program for local councils to prepare and implement CMPs.

Under the NSW Government's Coastal and Estuary Grants Program, funding has been given to relevant councils to commence preparation of CMPs for the Cooks River Catchment and Georges River Catchment. Stage 1, the scoping studies, has recently been completed and Stage 2 is progressing. The CMPs for these waterways and Botany Bay will identify catchment pressures and prioritise management options for issues relating to coastal and estuary health. Water quality management actions such as stormwater infrastructure improvements, restoring and maintaining riparian areas, and strategic land-use planning will be considered during the process.

Several Sydney councils implement the Georges River Estuary CZMP, including Liverpool, Sutherland Shire, Georges River and Bayside councils. With funding from the NSW Government's Coastal and Estuary Grants Program, a number of foreshore access areas or erosion hotspots have been identified for improvements to reduce sedimentation and improve water quality of the Georges River.

Randwick City Council

Randwick City Council operates and maintains 15 stormwater harvesting treatment systems with UV filtration across the local government area. These systems treat stormwater by removing suspended solids, bacteria and other organic and inorganic materials before it is used for irrigation in surrounding landscaped and garden areas, saving Randwick City Council approximately 455 megalitres of water (which equates to 187 Olympic sized swimming pools or \$1 million in cost savings).

Randwick City Council maintains 36 GPTs on stormwater lines leading to the local bays, which are all cleaned regularly. From February 2021 to February 2022, approximately 372 tonnes of material was removed from these GPTs. There is also a systematic cleaning program for all drainage pits including a regular street sweeping program that assists with reducing stormwater pollution to the local bays. Council continues to conduct litter education campaigns throughout the local government area to educate residents on the proper disposal of waste. This program aims to reduce the amount of litter disposed on beaches and entering the ocean.

Council officers undertake their routine inspections and regulatory duties to ensure stormwater pollution is investigated and mitigated to reduce impacts to the water quality of local recreational waterways.

Randwick City Council has a strategic program and reactive process to monitor and assess the condition of the stormwater pipes in the local area using CCTV.



Brighton Le Sands

Photo: Beachwatch/DPE

Bayside Council

Bayside Council continues to undertake water management related works and maintenance projects to maintain and improve water quality through its Water Management Strategy. This includes maintenance of aerators in waterbodies, ongoing maintenance of litter and sediment control traps preventing pollutants from entering the waterways, removal of aquatic weeds and excess sediment build-up in local waterways, education and Bushcare programs for the community as well as dune and estuarine vegetation restoration programs.

Georges River Council

Georges River Council continues to prevent litter, organic matter, sediment and oil entering local waterways through water sensitive urban design and GPT installation projects, including the Gannons Park Water Quality Improvement and Stormwater Harvesting Scheme. This large-scale landscape and stormwater treatment project focused on reinstating a section of Boggywell Creek to develop significant public space and improve the community's connection to environmental processes. This was completed by removing concrete stormwater infrastructure, and constructing/reinstating natural waterway features including swales, wetlands, ponds and bioretention systems. This project has been designed to remove 16,000 kg of sediment, 22 kg of phosphorus and 131 kg of nitrogen every year from stormwater, while providing a source of water for field irrigation.



Jew Fish Bay Baths

Photo: Beachwatch/DPE

During 2021–2022, council installed a GPT in Depot Creek in Mortdale to remove gross pollutants from entering Lime Kiln Bay, and completed remediation and naturalisation works at lower Poulton Park, redirecting stormwater through swales and detention points before it reaches the waterway.

Georges River Council has recently completed a wetland health assessment report for the Lime Kiln Bay constructed wetlands. This work was done to understand the current performance and ecological health of the wetland system.

Georges River Council has also endorsed the Georges River Foreshore Access and Improvement Plan. This plan represents a key component to deliver resilient estuarine ecosystems and vibrant communities along the foreshore areas of the Georges River within the Georges River Council local government area. The primary purpose is to enable

public land along the Georges River foreshore to be planned and managed for an optimal balance of environmental health and ecological resilience, and access, use and enjoyment by the community.

Sutherland Shire Council

Sutherland Shire Council continues to use the stormwater levy to fund programs in stormwater quality improvement and water quality monitoring. Council has installed 2 major GPTs on a trunk stormwater line draining parts of Taren Point, Caringbah and Miranda. Smaller GPTs have also been investigated for installation on multiple stormwater outlets to Gwawley Creek. Both initiatives will help improve water quality and amenity in the waterways of Sylvania Waters and the lower Georges River.

As part of its long-running Strategic Water Monitoring Program (SWaMP) council continues to monitor over 20 freshwater and estuarine waterways that flow into either the Georges River, Botany Bay or Bate Bay. The program tests physical, chemical and biological (macroinvertebrates) condition to measure the ecological health of our waterways.



Lilli Pilli Baths

Photo: Beachwatch/DPE

Council continues to rehabilitate the Shire's urban waterways, which in 2021–2022 included Loftus Creek, Carina Creek, Still Creek, Bottle Creek, Fahy Creek and Mandowie Creek. Rehabilitation works undertaken by specialist bush regeneration contractors include weed removal, creek bank stabilisation and revegetation with thousands of native plants grown at council's community nursery. These works improve local biodiversity and amenity, and contribute to improving the ecological health of receiving waterways.

For the past few years, council has actively supported the Georges Riverkeeper catchment management group in preparing a CMP for Georges River, culminating in 2021–2022 in the completion of the CMP Stage 1 Scoping Study. Water quality improvement through improved catchment management was identified as a key focus area for the CMP, and will be examined in more detail in the forthcoming CMP Stages 2–4.

Sydney Water

Sydney Water is implementing a major program of works and activities to improve the suitability for swimming at Foreshore Beach by reducing the occurrence of overflows to the Mill Stream during wet weather.

Sydney Water has inspected, cleaned and repaired sewer mains on the northern and western sides of Botany Bay that

have a high likelihood of discharging sewage to waterways if they become blocked. When significant tree root intrusion to the public sewer from the private sewer was identified, property owners were requested to remedy the problem.

Port Hacking

Sutherland Shire Council



Jibbon Beach

Photo: Beachwatch/DPE

Sutherland Shire Council continues to prioritise protecting the ecological health and recreational amenity of Port Hacking. On behalf of the NSW Government, council is project managing the dredging of the port's main navigation channels with dredged marine sand used to nourish the Bate Bay beaches.

In 2021–2022, council's Waterways Rehabilitation Program continued works at Coonong and Savilles creeks. Initial project works were commenced at an unnamed watercourse at Marina Crescent Reserve, Northwest Arm. Works included creek bank stabilisation, removal of woody weeds, exotic vines and scramblers, and revegetation of thousands of native plants grown at council's community nursery. The works will improve water quality, the area's aesthetics and biodiversity, and contribute to the condition of Port Hacking.

Council is currently preparing to commence a CMP for Port Hacking, starting with preparation of the Stage 1 Scoping Study. Navigation, waterway health and foreshore access will be some of the key focus areas for the CMP.

Sydney Water

Sydney Water has inspected, cleaned and repaired sewer mains on the northern side of Port Hacking that have a high likelihood of discharging sewage to waterways if they become blocked. When significant tree root intrusion to the public sewer from the private sewer was identified, property owners were asked to remedy the problem.

To reduce the incidence of wet weather sewage overflows in the catchments of Gunnamatta Bay Baths, Sydney Water has amplified pipes and pumps and included storage tanks across the Cronulla Peninsula.



Sampling sites at Beach Suitability Grades at Sydney's Southern beaches




Sampling sites and Beach Suitability Grades in Botany Bay and lower Georges River



Sampling sites and Beach Suitability Grades in Port Hacking

Boat Harbour

Beach grade: **G**



Boat Harbour is a 150 m long unpatrolled private beach at the northern end of Bate Bay. It is the beach closest to the Cronulla WWTP outfall at Potter Point.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater and onsite sewer systems behind the beach.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jul 2020 to Apr 2022	89%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Greenhills Beach

Beach grade: **VG**



Greenhills Beach is 3 km long and situated at the northern end of Bate Bay. The beach is not patrolled.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after 5 mm or more of rain.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Jun 2020 to Apr 2022	97%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Wanda Beach

Beach grade: **VG**



Wanda, Elouera and North Cronulla beaches form a 1.5 km stretch of beach towards the southern end of Bate Bay. Lifeguards patrol from October to April.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, but generally remained below the safe swimming limit across all rainfall categories.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Elouera Beach

Beach grade:



Wanda, Elouera and North Cronulla beaches form a 1.5 km stretch of beach towards the southern end of Bate Bay. Lifeguards patrol the beach from October to April.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit in response to 20 mm or more of rain.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality




Water quality in response to rainfall



North Cronulla Beach

Beach grade: **VG**



North Cronulla Beach is at the southern end of a 1.5 km stretch of beach in Bate Bay. Lifeguards patrol the beach all year round.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after rain.


See ‘How to read this report’ for key to map. The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



South Cronulla Beach

Beach grade:



South Cronulla beach is 300 m long and situated at the southern end of Bate Bay. Lifeguards patrol the beach all year round.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after rain.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	96%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Shelly Beach

Beach grade: **VG**



Shelly beach is 50 m long and is not patrolled by lifeguards. The adjacent ocean pool is the most suitable area for swimming.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit in response to 20 mm or more of rain.

See 'How to read this report' for key to map.


The site has been monitored since 1989.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	98%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality




Water quality in response to rainfall



Oak Park

Beach grade:



Oak Park beach is 15 m long, with the most suitable area for swimming adjacent to the ocean pool. Lifeguards do not patrol the swimming area.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit in response to 5 mm or more of rain.

The site has been monitored since 1989.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Ocean beach	Aug 2020 to Apr 2022	100%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: A



Dry and wet weather water quality



Water quality in response to rainfall



Silver Beach

Beach grade: **G**



Silver Beach is a netted swimming area at the centre of a 2.8 km long beach on the southern shore of Botany Bay.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	92%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Como Baths

Beach grade: **G**



Como Baths is approximately 25 m wide and backed by a narrow sandy beach in the lower Georges River.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream sources in the Georges River.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to no rain, and often after 5 mm or more.

The site has been monitored since 1994.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	87%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Jew Fish Bay Baths

Beach grade:



Jew Fish Bay Baths is a 200 m long netted swimming area located in Jew Fish Bay in the lower Georges River.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including upstream sources in the Georges River.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	90%	100	Declined


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality




Water quality in response to rainfall



Oatley Bay Baths



Beach grade: **P**



Oatley Bay Baths is a netted swimming area located on the western shore of Oatley Bay in the lower Georges River.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including upstream sources in the Georges River and stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	90%	100	Declined


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Carss Point Baths

Beach grade: P



Carss Point Baths is a netted swimming enclosure on the western shore of Kogarah Bay in the lower Georges River.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including upstream sources in the Georges River and stormwater.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after rainfall.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	87%	100	Declined ↓


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Sandringham Baths

Beach grade:



Sandringham Baths is a netted swimming area near the mouth of the Georges River and is backed by a small beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from the Georges River.


Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit after 5 mm or more of rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	99%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Dolls Point Baths

Beach grade: P



Dolls Point Baths is a netted swimming area with a sandy beach at the southern end of Lady Robinsons Beach in Botany Bay.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including discharge from the Georges River.

Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.


See 'How to read this report' for key to map. The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	77%	100	Declined ↓


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Ramsgate Baths

Beach grade:



Ramsgate Baths is a netted swimming enclosure with a sandy beach near the southern end of Lady Robinsons Beach in Botany Bay.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from the Georges River.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	90%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Monterey Baths

Beach grade:



Monterey Baths is a netted swimming area with a sandy beach located toward the southern end of Lady Robinsons Beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including discharge from the Georges River and stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Sep 2019 to Apr 2022	91%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Brighton-Le-Sands Baths

Beach grade:



Brighton-Le-Sands Baths is a netted swimming area at the centre of Lady Robinsons Beach in Botany Bay and is backed by a sandy beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including sewage overflows and river discharge.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	93%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Kyeemagh Baths

Beach grade:



Kyeemagh Baths is a netted swimming area with a sandy beach at the northern end of Lady Robinsons Beach, near the Cooks River mouth.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including the discharge from the Cooks River and sewage overflows.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	86%	100	Declined


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Foreshores Beach

Beach grade:



Foreshores Beach is an unnetted sandy beach in Botany Bay. It is located near a boat ramp, and is adjacent to the Sydney Airport runway and Port Botany.

The Beach Suitability Grade of Very Poor indicates microbial water quality is very susceptible to faecal pollution with many potential sources of faecal contamination including stormwater and sewage overflows that discharge into Mill Stream.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after rainfall.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	77%	100	Declined


Sanitary inspection: High




Microbial Assessment Category: D



Dry and wet weather water quality



Water quality in response to rainfall



Yarra Bay

Beach grade:



Yarra Bay is a 750 m long sandy beach in Botany Bay. The swimming area is not netted and has a rock groyne at the southern end.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with several potential sources of faecal contamination including stormwater that ponds in the middle of the beach.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after rainfall.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	92%	100	Declined


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Frenchmans Bay



Beach grade:



Frenchmans Bay is a 500 m long sandy beach in Botany Bay. The swimming area is not netted.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with potential faecal contamination from several sources including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	77%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: C



Dry and wet weather water quality



Water quality in response to rainfall



Congwong Bay

Beach grade:



Congwong Bay is a 150 m long beach near the mouth of Botany Bay. The swimming area is not netted.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of minor faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Nov 2019 to Apr 2022	88%	100	Stable


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Jibbon Beach

Beach grade:



Jibbon Beach is located at the entrance to Port Hacking. The beach is backed by the Royal National Park and accessed from Bundeena.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination.


Enterococci levels increased slightly with rainfall, occasionally exceeding the safe swimming limit in response to light rain, and often after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1999.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Jun 2019 to Apr 2022	95%	100	Declined


Sanitary inspection: Low




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Horderns Beach

Beach grade:



Horderns Beach is located on the southern shore of Port Hacking and is backed by the town of Bundeena.



The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including Bundeena Creek.


Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and regularly after 20 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1999.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Jun 2019 to Apr 2022	92%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



Gymea Bay Baths

Beach grade: P



Gymea Bay Baths is an enclosed tidal swimming area backed by a narrow sandy beach in the upper reaches of Port Hacking.

The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with potential faecal contamination from several sources including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after no rain, and often after rainfall.

See 'How to read this report' for key to map.


The site has been monitored since 1999.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Aug 2019 to Apr 2022	82%	100	Stable


Sanitary inspection: High




Microbial Assessment Category: C



Dry and wet weather water quality




Water quality in response to rainfall



Lilli Pilli Baths

Beach grade:



Lilli Pilli Baths is a netted tidal swimming area on the western side of Lilli Pilli Point in Port Hacking.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 5 mm or more.

The site has been monitored since 1999.


See 'How to read this report' for key to map.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	92%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality




Water quality in response to rainfall



Gunnamatta Bay Baths

Beach grade:



Gunnamatta Bay Baths is a netted tidal swimming area in the lower reaches of Port Hacking and is backed by a narrow beach.

The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.


Enterococci levels increased with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 5 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 1994.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status
Estuarine	Oct 2019 to Apr 2022	89%	100	Stable


Sanitary inspection: Moderate




Microbial Assessment Category: B



Dry and wet weather water quality



Water quality in response to rainfall



How to read this report

Beach Suitability Grades

Beach Suitability Grades provide an assessment of the suitability of a swimming location for recreation over time and are based on a combination of sanitary inspection (identification and rating of potential pollution sources at a beach) and microbial assessment (water quality measurements gathered over previous years). There are 5 grades ranging from Very Good to Very Poor:

VG Very Good

Location has generally excellent microbial water quality and very few potential sources of faecal pollution. Water is considered suitable for swimming almost all of the time

G Good

Location has generally good microbial water quality and water is considered suitable for swimming most of the time. Swimming should be avoided during and for up to one day following heavy rain at ocean beaches and up to 3 days at estuarine sites

F Fair

Microbial water quality is generally suitable for swimming, but because of the presence of significant sources of faecal contamination, extra care should be taken to avoid swimming during and for up to 3 days following rainfall or if there are signs of pollution such as discoloured water or odour or debris in the water

P Poor

Location is susceptible to faecal pollution and microbial water quality is not always suitable for swimming. During dry weather conditions, ensure that the swimming location is free of signs of pollution, such as discoloured water, odour or debris in the water, and avoid swimming at all times during and for up to 3 days following rainfall

VP Very Poor

Location is very susceptible to faecal pollution and microbial water quality may often be unsuitable for swimming. It is generally recommended to avoid swimming at these sites almost all of the time

Some of the Beach Suitability Grades in this report are **provisional**, as the information required for the analysis is incomplete due to limited bacterial data or limited information on potential pollution sources in a beach catchment.

The guidelines

The National Health and Medical Research Council's guidelines for managing risks in recreational water (NHMRC 2008) were adopted for use in NSW in May 2009. These guidelines have been adopted in all Australian states and territories and are supported by guidance notes developed by the Department of Health Western Australia (WA Department of Health 2007).

Enterococci

The national guidelines advocate the use of enterococci as the single preferred faecal indicator in marine waters.

These bacteria are excreted in faeces and are rarely present in unpolluted waters. Enterococci have shown a clear dose–response relationship to disease outcomes in marine waters in the northern hemisphere. In accordance with the guidelines, Beachwatch tests for enterococci only. The enterococci density in water samples is analysed in the laboratory using method AS/NZS 4276.9:2007 (Standards Australia 2007).

Enterococci are measured in colony forming units per 100 mL of sample (cfu/100 mL).

Beach Suitability Grades are determined by using the following matrix:

		Microbial Assessment Category			
		A	B	C	D
Sanitary Inspection Category	Very Low	Very Good	Very Good	Follow Up	Follow Up
	Low	Very Good	Good	Follow Up	Follow Up
	Moderate	Good	Good	Poor	Poor
	High	Good	Fair	Poor	Very Poor
	Very High	Follow Up	Fair	Poor	Very Poor

Using the Beach Suitability Grade classification matrix, sites assigned a moderate Sanitary Inspection Category can only be rated as Good or Poor, with no option of Fair grades. This can create the impression of a large change in water quality when in fact there need only be a slight increase in bacterial counts to push it over the threshold, with no significant increase in the risk to public health.

Microbial Assessment Category (MAC)

There are 4 Microbial Assessment Categories (A to D) and these are determined from the 95th percentile of an enterococci dataset of at least 100 data points. Each MAC is associated with a risk of illness determined from epidemiological studies. The risks of illness shown below are not those associated with a single data point but are the overall risk of illness associated with an enterococci dataset with that 95th percentile (Wyer et al. 1999).

Risk of illness associated with Microbial Assessment Categories

Category	Enterococci (cfu/100 mL)	Illness risk*
A	≤40	GI illness risk: <1% AFR illness risk: <0.3%
B	41–200	GI illness risk: 1–5% AFR illness risk: 0.3–1.9%
C	201–500	GI illness risk: >5–10% AFR illness risk: >1.9–3.9%
D	>500	GI illness risk: >10% AFR illness risk: >3.9%

* GI = gastrointestinal illness; AFR = acute fever and rash

Calculating the MAC

The 95th percentile is a useful statistic for summarising the distribution of enterococci data at a site. It embodies elements of both the location of the distribution (how high/low the enterococci counts are) and the scale of the distribution (how variable the enterococci counts are).

The 95th percentile values for each of the 4 Microbial Assessment Categories were determined by the World Health Organization using enterococci data collected from swimming locations across Europe. These values will represent different probabilities of illness if the distribution of enterococci data from swimming locations in NSW differs from the European distribution.

In recognition of this issue, Dr Richard Lugg (Department of Health, Western Australia) has developed a Microsoft® Excel tool for calculating a modified 95th percentile that takes into account the distribution of data. This tool has been used to calculate the 95th percentile values presented in this report and has been adopted for use by other state governments in Australia.

The tool can be downloaded from the WA Government's 'Environmental waters publications' webpage, under *Forms and templates*.

Sanitary Inspection Category (SIC)

More information about the **sanitary inspection** process is available on the DPE 'Sanitary inspection of beaches' webpage.

The aim of a sanitary inspection is to identify all sources of faecal contamination that could affect a swimming location and assess the risk to public health posed by these sources. It is an assessment of the likelihood of bacterial contamination from identified pollution sources and should, to some degree, correlate with the bacterial water quality results obtained from sampling.

The main sources of faecal contamination considered in the sanitary inspection are: bathers, toilet facilities, wastewater treatment plants (WWTPs), sewage overflows, sewer chokes, onsite systems, wastewater re-use, stormwater, river discharge, lagoons, boats and animals.

Rivers, lakes and estuaries themselves can be potential sources of faecal contamination to sites located in these waterbodies, with contaminated water from upstream or surrounding areas impacting water quality at the swimming location. This source is captured in river discharge or lagoon category, and shown as the waterbody in the sanitary inspection charts.

Through the sanitary inspection process, beaches are categorised to reflect the overall likelihood of faecal contamination. There are 5 categories: Very Low, Low, Moderate, High and Very High.



Stormwater drain flow
Photo: Beachwatch/DPE

Stormwater in urban areas often contains sewage from leakages, overflows or sewer chokes when the sewerage system fails.

Sewage overflows can occur in wet weather when the network has exceeded capacity due to rainwater entering the system. The mix of sewage and rainwater discharges from designated overflow points and drains to waterways, usually via the stormwater system. Overflows from the sewerage system can also occur in dry weather due to mechanical failure or power outage.

Sewer chokes occur due to blockages in the pipes usually due to tree roots, oil, grease or debris. This causes sewage to back up and escape via sewer inspection points, designed overflow structures or cracks in the pipes, then drain to waterways, usually via the stormwater system.

Explanation of tables

Each region contains tables listing all monitored swimming sites including site type, beach grade and change in grade from the previous year.

The following symbols are used to show the change in beach grade from the previous year:


- Stable
- ↑ Improved
- ↓ Declined

A provisional grade indicates the assessment is based on limited data collected during the assessment period and should not be compared to the beach grade from the previous year.

Explanation of graphs, charts, and information bars on beach pages

Microbial Assessment Category (MAC) chart


On each beach page, the MACs for the last 5 years are displayed on a simple bar chart. The MAC for the current year is based on enterococci data collected during the assessment period. The bars are labelled with the 95th percentile value for each year and the thresholds dividing the A, B, C and D categories are marked in green, amber and red for reference.



Sanitary Inspection Category (SIC) chart


The results of the sanitary inspection for each swimming location are presented in a radar pie chart. The chart shows the likelihood that each identified pollution source will contribute to faecal contamination at a swimming site, as indicated by the size and colour of the segment, ranging from

very low (lightest colour) to very high (darkest colour) as shown below. The sum of these contributions is the overall likelihood, or Sanitary Inspection Category.



Wet and dry weather water quality chart


Enterococci levels in wet and dry weather conditions are presented for each swimming location as a bar graph. All data collected during the assessment period is included in the analysis. Dry weather is defined as no rainfall recorded in the previous 24 hours. Each bar is colour coded to show the number of enterococci results up to 40 cfu/100 mL, between 41 and 200 cfu/100 mL, between 201 and 500 cfu/100 mL and greater than 500 cfu/100 mL. These categories reflect the Microbial Assessment Category thresholds and are coloured on the graph as dark green, light green, amber and red respectively.




It is expected that swimming sites with lower levels of flushing will show some elevated bacterial results in dry weather samples (no rainfall in the previous 24 hours) due to the longer time needed to recover from a rainfall event. At some estuarine and lake/lagoon swimming locations the impacts of stormwater pollution on beach water quality may be detected up to 3 days after rainfall.

Water quality in response to rainfall

Trends in enterococci levels in response to rainfall are shown using a box plot. For reference, enterococci levels of 40 cfu/100 mL and 200 cfu/100 mL are indicated with a green and orange line, respectively. The 40 cfu/100 mL level is referred to as the 'safe swimming limit'. The enterococci data were obtained from the last 5 years of monitoring. Rainfall data were obtained from rain gauges situated close to the sample site and are 24-hour totals to 9am on the day of sampling. If there are fewer than 5 enterococci data points in a rainfall category, individual data points are presented instead of a box plot. At sites where many results are below the detection limit (1 cfu/100 mL), only the upper portion of the box plots will be visible.



Each part of the box plot represents a significant percentile value of the sample population:



Information bars
















Information bars on each beach page provide a summary of details about the swimming site.

The **assessment period** shows the timeframe in which the water samples were collected. The NHMRC guidelines state beach grades should be determined from the most recent 100 water quality results collected within a 5-year period. The assessment period varies between sites depending on sampling frequency.

Dry weather samples suitable for swimming (**dry weather swimmability**) shows the percentage of water samples with enterococci levels below 40 cfu/100 mL. Dry weather is defined as no rainfall in the previous 24 hours. Swimming sites with lower levels of flushing often have a lower percentage of dry weather samples within the safe swimming limit due to the impacts of rainfall detected up to 3 days after the event.

Explanation of maps

A map of individual swimming locations is presented on each beach page. The scale of the maps is 1:10,000. Each map shows the location of the sampling site, land use and features such as surf lifesaving clubs. Potential pollution sources such as stormwater drains, sewage pumping stations, wastewater treatment plants, lagoons, rivers and creeks, are shown where accurate data is held.

Key to maps	
	Sampling Site
	Surf Life Saving Club
	Wastewater Treatment Plant
	Sewage Pumping Station
	Sewage Overflow
	Stormwater Drain
	Water
	Baths
	National Park/Reserve/ Other Park
	Built-up Area
	Sand
	Roads
	Major Roads
	Baths – Netted Area
	Breakwater/Wharf

References

NHMRC (2008) *Guidelines for managing risks in recreational water*, National Health and Medical Research Council, Australian Government Publishing Service, Canberra, ACT.

Standards Australia (2007) *AS/NZS 4276.9:2007, Water microbiology Method 9: Enterococci – Membrane filtration method (ISO 7899-2:2000, MOD)*, Standards Australia International Ltd, Sydney and Standards New Zealand, Wellington.

WA Department of Health (2007), *Microbial quality of recreational water guidance notes in support of chapter 5 of the National Health and Medical Research Council guidelines for managing risks in recreational water, 2006*, Department of Health, Western Australia and The University of Western Australia, October 2007, ww2.health.wa.gov.au/Articles/A_E/Environmental-waters-publications, accessed 23/06/22.

Wyer MD, Kay D, Fleisher JM, Salmon RL, Jones F, Godfree AF, Jackson G and Rogers A (1999) 'An experimental health related classification for marine waters', *Water Research*, 33(3):715–722.

More information

- [Beachwatch NSW on Twitter](#)
- [Beachwatch NSW on Facebook](#)
- [Beachwatch webpage](#)
- [Coastal management program progress](#)
- [Sanitary inspection of beaches](#)
- [Subscribe to environment and heritage newsletters](#)
- [Towards Safer Swimming: Rose Bay](#)
- [WA Government environmental water publications](#)