

Beachwatch

# State of the beaches 2023–24

South Coast region

Department of Climate Change, Energy, the Environment and Water



## Acknowledgement of Country

Department of Climate Change, Energy, the Environment and Water acknowledges the Traditional Custodians of the lands where we work and live.

We pay our respects to Elders past, present and emerging.

This resource may contain images or names of deceased persons in photographs or historical content. © 2024 State of NSW and Department of Climate Change, Energy, the Environment and Water

With the exception of photographs, the State of NSW and Department of Department of Climate Change, Energy, the Environment and Water (the department) are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required to reproduce photographs.

Learn more about our copyright and disclaimer at www.environment.nsw.gov.au/copyright

Artist and designer Nikita Ridgeway from Aboriginal design agency Boss Lady Creative Designs created the People and Community symbol.

Cover photo: Narooma Main Beach, Eurobodalla. Beachwatch/DCCEEW

Published by: Environment and Heritage Department of Climate Change, Energy, the Environment and Water Locked Bag 5022, Parramatta NSW 2124 Phone: +61 2 9995 5000 (switchboard) Phone: 1300 361 967 (Environment and Heritage enquiries) TTY users: phone 133 677, then ask for 1300 361 967 Speak and listen users: phone 1300 555 727, then ask for 1300 361 967 Email <u>info@environment.nsw.gov.au</u>

Website www.environment.nsw.gov.au

ISBN 978-1-923285-38-5 EH 2024/0256 September 2024

#### Find out more about your environment at:

#### environment.nsw.gov.au

# Contents

South Coast region summary 2023–2024	1
Monitoring water quality for swimming in New South Wales	1
Rainfall impacts	2
Shoalhaven City Council	7
Overall results	7
Shoalhaven Heads Beach	10
Tilbury Cove	11
Warrain Beach	12
Collingwood Beach	13
Cudmirrah Beach	14
Mollymook Beach	15
Rennies Beach	16
Racecourse Beach	17
Bawley Point Beach	18
Merry Beach	19
Eurobodalla Shire Council	20
Overall results	20
Cookies Beach	25
Caseys Beach	26
Surf Beach	27
Malua Bay Beach	28
Broulee Beach	29
South Broulee (Bengello) Beach	30
Shelley Beach	31
Tuross Main Beach	32
Brou Beach	33
Wagonga Inlet	34
Narooma Main Beach	35

How to read this report	36
Beach Suitability Grades	36
Explanation of tables	42
Explanation of graphs, charts, and information bars on beach pages	43
References	47
More information	47

Recreational water quality has been monitored in the South Coast region since 2002 by Shoalhaven City Council and Eurobodalla Shire Council under the Department of Climate Change, Energy, the Environment and Water's Beachwatch Partnership Program. This report summarises the performance of 21 swimming sites on the south coast of New South Wales, providing a long-term assessment of how suitable a site is for swimming. Monitored sites include ocean beaches and an estuarine swimming site in Wagonga Inlet.

In 2023–2024, 95% of swimming sites in the South Coast region were graded as Good or Very Good, including 19 ocean beaches. These sites were suitable for swimming for most or almost all of the time. This is an excellent result, and a similar performance to the previous year.



Mollymook Beach Photo: Beachwatch/DCCEEW

See the section on **Quality assurance** in the Statewide Summary for results of the quality assurance program.

# South Coast region summary 2023–2024

# Monitoring water quality for swimming in New South Wales

The water quality of beaches and other swimming locations is monitored under the NSW Government's Beachwatch programs to provide the community with accurate information on the cleanliness of the water and to enable individuals to make informed decisions about where and when to swim. Routine assessment also measures the impact of pollution sources, enables the effectiveness of stormwater and wastewater management practices to be assessed and highlights areas where further work is needed.

Swimming sites in New South Wales are graded as Very Good, Good, Fair, Poor or Very Poor in accordance with the National Health and Medical Research Council's 2008 *Guidelines for Managing Risks in Recreational Waters*. These Beach Suitability Grades provide a long-term assessment of how suitable a beach is for swimming. The grades are determined from the most recent 100 water quality results (2–4 years' worth of data depending on the sampling frequency) and a risk assessment of potential pollution sources.

Recreational water quality has been monitored on the South Coast by Shoalhaven Council and Eurobodalla Shire Council since 2002.

A **quality assurance** program ensures the information collected and reported by Beachwatch and its partners is accurate and reliable.

# **Rainfall impacts**

During 2023–2024, 21 swimming sites were monitored including ocean beaches and an estuarine swimming site. Rainfall is the major driver of pollution to recreational waters, generating stormwater runoff and triggering untreated discharges from the wastewater treatment and transport systems. Changes in rainfall patterns are reflected in beach water quality over time due to variation in the frequency and extent of stormwater and wastewater inputs.

The Beach Suitability Grades for 2023–2024 are based on water quality data collected over the last 2–4 years. Rainfall over this period has been diverse:


- 2020–2021: variable rainfall with some wet months
- 2021–2022: a very wet summer and autumn, including significant wet weather and flooding events
- 2022–2023: varied rainfall, with some very wet months over winter and spring, including the wettest July and October on record
- 2023–2024: mostly average to below average rainfall, except for some isolated wet months.


Rainfall on the South Coast was average to below average for the 2023–2024 reporting year, except for some isolated wet months. Well above average rainfall was recorded in November and December 2023 and April 2024 across the region.


November and December 2023 were notably wet on the South Coast. Jervis Bay and Ulladulla both recorded their highest November daily rainfall and record highest November and December total rainfall. Similarly, Narooma recorded its highest November total rainfall in 23 years.

See the section on How to read this report on page 36 for an explanation of the graphs, tables and Beach Suitability Grades.

#### South Coast region rainfall







3

#### **Health risks**

Contamination of recreational waters with faecal material from animal and human sources can pose significant health problems to beach users owing to the presence of pathogens (disease-causing microorganisms) in the faecal material. The most common groups of pathogens found in recreational waters are bacteria, protozoans and viruses.

Exposure to contaminated water can cause gastroenteritis, with symptoms including vomiting, diarrhoea, stomach-ache, nausea, headache and fever. Eye, ear, skin and upper respiratory tract infections can also be contracted when pathogens come into contact with small breaks and tears in the skin or ruptures of the delicate membranes in the ear or nose.

Certain groups of users may be more vulnerable to microbial infection than others. Children, the elderly, people with compromised immune systems, tourists, and people from culturally and linguistically diverse backgrounds are generally most at risk.

#### Beach Suitability Grades for South Coast region

Swimming site	Site type	Beach Suitability Grade	Change
Shoalhaven City Council			
Shoalhaven Heads Beach	Ocean beach	VG	$\bigcirc$
Tilbury Cove	Ocean beach	VG	$\bigcirc$
Warrain Beach	Ocean beach	VG	$\bigcirc$
Collingwood Beach	Ocean beach	VG	$\bigcirc$
Cudmirrah Beach	Ocean beach	VG	$\bigcirc$
Mollymook Beach	Ocean beach	VG	$\bigcirc$
Rennies Beach	Ocean beach	VG	$\bigcirc$
Racecourse Beach	Ocean beach	VG	$\bigcirc$
Bawley Point Beach	Ocean beach	VG	$\bigcirc$
Merry Beach	Ocean beach	VG	$\bigcirc$
Eurobodalla Shire Council			
Cookies Beach	Ocean beach	VG	$\bigcirc$
Caseys Beach	Ocean beach	G	$\bigcirc$
Surf Beach	Ocean beach	P	0
Malua Bay Beach	Ocean beach	VG	$\bigcirc$
Broulee Beach	Ocean beach	VG	•
South Broulee (Bengello) Beach	Ocean beach	G	+
Shelley Beach (Moruya Heads)	Ocean beach	G	$\bigcirc$
Tuross Main Beach	Ocean beach	G	$\bigcirc$
Brou Beach	Ocean beach	VG	$\bigcirc$
Wagonga Inlet	Estuarine	G	$\bigcirc$
Narooma Main Beach	Ocean beach	G	$\bigcirc$

Beach Suitability Grade					•	Change	
VG	G	F	Р	VP		$\bigcirc$	+
Very Good	Good	Fair	Poor	Very Poor	Improved	Stable	Declined

# Shoalhaven City Council

# **Overall results**

All 10 swimming sites were graded as Very Good in 2023–2024. This is an outstanding result and a similar performance to previous years.

#### Percentage of sites graded as Very Good or Good

		2022- 2023		Trend
Ocean beaches (10 sites)	100%	100%	100%	

Ten swimming sites were monitored by Shoalhaven City Council. Samples were collected weekly between December and February and sampling and laboratory analysis was fully funded by the council.

100%

ocean beaches

graded Very

Good

See the section on **How to read this report** on page 36 for an explanation of the graphs, tables and Beach Suitability Grades.

#### **Best beaches**

Shoalhaven Heads Beach, Tilbury Cove, Warrain Beach, Collingwood Beach, Cudmirrah Beach, Mollymook Beach, Rennies Beach, Racecourse Beach, Bawley Point Beach and Merry Beach.

These sites had excellent water quality and were suitable for swimming almost all of the time.



Site types in Shoalhaven City Council

Ocean beaches were the only site type monitored in the Shoalhaven region.

As a general precaution swimming should be avoided during and for at least one day after heavy rain at ocean beaches, or if there are signs of stormwater pollution such as discoloured water or floating debris.




Beach Suitability Grades for Shoalhaven City Council ocean beaches

## **Ocean beaches**

All 10 ocean beaches continued to be graded as Very Good in 2023–2024: Shoalhaven Heads Beach, Tilbury Cove, Warrain Beach, Collingwood Beach, Cudmirrah Beach, Mollymook Beach, Rennies Beach, Racecourse Beach, Bawley Point Beach and Merry Beach.

While water quality at these sites was suitable for swimming almost all of the time, elevated bacterial levels were occasionally recorded at some of the beaches following heavy rainfall.





Sampling sites and Beach Suitability Grades in Shoalhaven City Council

# Shoalhaven Heads Beach





Shoalhaven Heads Beach is located towards the southern end of Seven Mile Beach at Shoalhaven Heads. The beach is patrolled over the summer months.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels had little response to rainfall and generally remained below the safe swimming limit across all rainfall categories.

See 'How to read this report' for key to map.


The site was monitored from 2003 to 2004 and since 2006.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	97%	55	Stable	$\bigcirc$

#### **Sanitary inspection: Low**





#### Dry and wet weather water quality



#### <=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL


#### **Microbial Assessment Category: A**





# **Tilbury Cove**





Tilbury Cove is located towards the south-eastern corner of Culburra Beach.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few sources of minor faecal contamination.


Enterococci levels had little response to rainfall and remained below the safe swimming limit across all rainfall categories.

The site was monitored from 2002 to 2004 and since 2006.


See 'How to read this report' for key to map.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	100%	57	Stable	$\bigcirc$

#### Sanitary inspection: Low





# Dry and wet weather water quality



#### 600 D

**Microbial Assessment Category: A** 





# Warrain Beach





Warrain Beach is located to the south of Penguin Head. The beach is patrolled over the summer months.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall but generally remained below the safe swimming limit across all rainfall categories.

See 'How to read this report' for key to map.


The site has been monitored since 2007.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	100%	56	Stable	$\bigcirc$


#### Sanitary inspection: Low




#### **Microbial Assessment Category: A**



#### Dry and wet weather water quality





# **Collingwood Beach**





Collingwood Beach is located in Jervis Bay, adjacent to the town of Vincentia. The beach is approximately 2 km long.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall but remained below the safe swimming limit across all rainfall categories.

See 'How to read this report' for key to map.


The site has been monitored since 2006.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	100%	57	Stable	$\bigcirc$

#### **Sanitary inspection: Low**





#### Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL

#### **Microbial Assessment Category: A**





# Cudmirrah Beach





Cudmirrah Beach is the main surf beach for the township of Sussex Inlet. The beach is approximately 3 km long and is patrolled over the summer months.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall but generally remained below the safe swimming limit across all rainfall categories.

See 'How to read this report' for key to map.


The site was monitored from 2003 to 2004 and since 2006.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grad status	de
Ocean beach	Dec 2019 to Feb 2024	100%	56	Stable	$\bigcirc$

#### Sanitary inspection: Low





#### Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL

#### **Microbial Assessment Category: A**





# Mollymook Beach





See 'How to read this report' for key to map.

Mollymook Beach is a popular beach that stretches for approximately 2 km. The beach is patrolled during the summer months.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall and generally remained below the safe swimming limit across most rainfall categories.


The site was monitored from 2002 to 2003 and since 2008.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	100%	57	Stable	$\bigcirc$

#### Sanitary inspection: Low





#### Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL

#### **Microbial Assessment Category: A**





# **Rennies Beach**





Rennies Beach is located near the town of Ulladulla. The beach is approximately 600 m long.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.


Enterococci levels had little response to rainfall and remained below the safe swimming limit across all rainfall categories.

The site has been monitored since 2006.


See 'How to read this report' for key to map.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	100%	56	Stable	$\bigcirc$

#### Sanitary inspection: Low





#### Dry and wet weather water quality



■<=40 cfu/100mL ■41-200 cfu/100mL ■ 201-500 cfu/100mL ■>500 cfu/100mL

#### **Microbial Assessment Category: A**





# **Racecourse Beach**





Racecourse Beach is located near the town of Ulladulla. The beach is approximately 1 km long.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall but generally remained below the safe swimming limit.

See 'How to read this report' for key to map.


The site was monitored from 2002 to 2004 and since 2006.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach grade status	
Ocean beach	Dec 2019 to Feb 2024	100%	57	Stable	)


#### Sanitary inspection: Low




#### Dry and wet weather water quality



#### **Microbial Assessment Category: A**





# **Bawley Point Beach**





See 'How to read this report' for key to map.

Bawley Point Beach is located on the northern side of Bawley Point and is approximately 250 m long.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.

Enterococci levels had little response to rainfall and generally remained below the safe swimming limit across all rainfall categories.


The site has been monitored since 2006.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	96%	56	Stable	$\bigcirc$


#### Sanitary inspection: Low




#### Dry and wet weather water quality



#### **Microbial Assessment Category: A**





# Merry Beach





See 'How to read this report' for key to map.

Merry Beach is located south of the town of Kioloa. The beach is approximately 400 m long and is backed by a reserve and caravan park.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after 20 mm or more of rainfall.


The site has been monitored since 2006.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Dec 2019 to Feb 2024	96%	56	Stable	$\bigcirc$


#### Sanitary inspection: Low




#### Dry and wet weather water quality



#### **Microbial Assessment Category: A**





# Eurobodalla Shire Council

# **Overall results**

Ten of the 11 swimming sites were graded as Very Good or Good in 2023–2024. This is a similar performance to the previous year.

#### Percentage of sites graded as Very Good or Good

	2021– 2022	2022- 2023	2023- 2024	Trend
Ocean beaches (10 sites)	80%	90%	90%	
Estuarine sites (1 sites)	100%	100%	100%	

Eleven swimming locations were monitored by Eurobodalla Shire Council. Samples were collected weekly between November and March and sampling and analysis was fully funded by the council.

90%

ocean beaches

graded Good or Very Good

> See the section on How to read this report on page 36 for an explanation of the graphs, tables and Beach Suitability Grades.

#### **Best beaches**

Cookies Beach, Malua Bay Beach, Broulee Beach and Brou Beach.

These sites had excellent water quality and were suitable for swimming almost all of the time.

20



Site types in Eurobodalla Shire Council

Swimming sites monitored in the Eurobodalla region include ocean beaches and an estuarine area in Wagonga Inlet, with each site type having a different response to rainfall-related impacts.

In general, estuarine swimming sites do not perform as well as ocean beaches, due to lower levels of flushing increasing the time needed to disperse and dilute pollution inputs, taking longer to recover from stormwater events.

As a general precaution swimming should be avoided during and for at least one day after heavy rain at ocean beaches, and for up to 3 days in estuarine areas, or if there are signs of stormwater pollution such as discoloured water or floating debris.

#### **Ocean beaches**

Nine of the 10 ocean beaches were graded as Very Good or Good in 2023–2024.

Cookies Beach, Malua Bay Beach, Broulee Beach and Brou Beach were graded as Very Good in 2023–2024. Water quality at these sites was suitable for swimming almost all of the time.

Caseys Beach, South Broulee (Bengello) Beach, Shelley Beach, Tuross Main Beach and Narooma Main Beach were graded as Good in 2023–2024. Water quality at these sites was frequently suitable for swimming during dry weather conditions, with between 90% and 97% of dry weather samples within the safe swimming limit. However, enterococci levels occasionally exceeded the safe swimming limit after little or no rain. South Broulee (Bengello) Beach declined to Good from Very Good in the previous year due to a slight decline in microbial water quality in the assessment period.

Surf Beach was graded as Poor in 2023–2024, consistent with the previous years. Elevated bacterial levels were often recorded during dry weather conditions and increased with increasing rainfall. Despite this, 77% of dry weather samples were within the safe swimming limit.

Microbial water quality at Surf Beach has improved slightly this year, despite a higher proportion of wet weather samples in this year's assessment period compared to the previous year's assessment period. Microbial water quality at this site had been declining for several years prior to this result.

Council has been investigating the water quality at Surf Beach since 2019, which at times is identified as poor. Council undertook an extensive monitoring program to investigate potential sources of contamination in the



Beach Suitability Grades for Eurobodalla Shire Council ocean beaches

catchments draining to Surf Beach Creek and Surf Beach. Following this, council engaged a consultant to undertake a water quality investigation that included statistical analysis, hydrodynamic modelling, investigative sampling and DNA marker analysis. The investigation identified multiple potential sources of contamination including domestic canine faeces entering waterways, discharge from Batemans Bay Sewage Treatment Plant (STP) during overflow events. STP outfall contamination via tidal patterns and failure or leakage of onsite sewer management systems (OSSM) upstream of Surf Beach. Council will action the recommendations to improve water quality, including a responsible dog waste disposal education program and an audit of all OSSMs in the catchment to reduce any potential contamination from these sources. Council also completed the Surf Beach STP upgrades in December 2023.

It is recommended to avoid swimming during and for at least one day following rainfall at ocean beaches or if there are signs of stormwater pollution such as discoloured water and floating debris.



Beach Suitability Grades for Eurobodalla Shire Council estuarine beaches

## **Estuarine beaches**

Wagonga Inlet continued to be graded as Good in 2023– 2024, similar to previous years. While water quality at this site was mostly suitable for swimming, elevated enterococci results were occasionally recorded after light rain, and often after moderate rainfall. Pollution inputs from elsewhere within Wagonga Inlet may impact water quality at this site.

Swimming should be avoided during and for up to 3 days after rainfall, or if there are signs of pollution such as discoloured water or floating debris.



Sampling sites and Beach Suitability Grades in Eurobodalla Shire Council

# **Cookies Beach**





Cookies Beach is located near the town of South Durras. Murramarang National Park lies to the south.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit after 20 mm or more of rain.


See 'How to read this report' for key to map. The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade
type	period		samples	status
Ocean beach	Nov 2020 to Mar 2024	94%	100	Stable


600

95th %ile


#### Sanitary inspection: Low



#### Dry and wet weather water quality



32



#### Microbial Assessment Category: A

32

2019-20 2020-21 2021-22 2022-23 2023-24

Water quality in response to rainfall

Microbial Assessment Category

# **Caseys Beach**





See 'How to read this report' for key to map.

Caseys Beach is approximately 800 m long and is located to the south of Observation Point.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including river discharge.

Enterococci levels generally increased with increasing rainfall, occasionally exceeding the safe swimming limit in response to little or no rain, and often after 10 mm or more.


The site has been monitored since 2002.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach gra status	de
Ocean beach	Nov 2020 to Mar 2024	90%	100	Stable	$\bigcirc$

#### **Sanitary inspection: Moderate**





#### Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL = 201-500 cfu/100mL =>500 cfu/100mL

#### **Microbial Assessment Category: B**





# Surf Beach





See 'How to read this report' for key to map.

Surf Beach is a popular beach approximately 350 m long and is patrolled in the warmer months.


The Beach Suitability Grade of Poor indicates microbial water quality is susceptible to faecal pollution, particularly after rainfall and occasionally during dry weather conditions, with potential faecal contamination from stormwater.

Enterococci levels increased with increasing rainfall, often exceeding the safe swimming limit after little or no rain, and regularly after 10 mm or more.


The site has been monitored since 2002.

Site type	Assessment period	Dry weather samples suitable for swimming	Water samples	Beach gra status	de
Ocean beach	Nov 2020 to Mar 2024	77%	100	Stable	$\bigcirc$


#### **Sanitary inspection: Moderate**




#### Dry and wet weather water quality



Microbial Assessment Category: D





# Malua Bay Beach





Malua Bay Beach is approximately 500 m long and is patrolled during the warmer months.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.

Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit in response to 20 mm or more of rain.


See 'How to read this report' for key to map. The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade
type	period		samples	status
Ocean beach	Nov 2020 to Mar 2024	96%	100	Stable


#### Sanitary inspection: Low




#### **Microbial Assessment Category: A**



#### Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL



# **Broulee Beach**





Broulee Beach extends from Candlagan Creek in the north to Broulee Island in the south.

The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few significant sources of faecal contamination.


Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain and often after 10 mm or more.

See 'How to read this report' for key to map.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Nov 2020 to Mar 2024	93%	100	Improved	


#### Sanitary inspection: Low




#### Dry and wet weather water quality



#### **Microbial Assessment Category: B**





# South Broulee (Bengello) Beach





See 'How to read this report' for key to map.

Bengello Beach extends from Broulee Head to the mouth of the Moruya River. The beach is patrolled during the summer months.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including stormwater.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after light rain, and often after 10 mm or more.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade
type	period		samples	status
Ocean beach	Nov 2020 to Mar 2024	93%	100	Declined

#### Sanitary inspection: Low





Dry and wet weather water quality



<=40 cfu/100mL =41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL

#### **Microbial Assessment Category: A**





# Shelley Beach

# TASMAN SEA

See 'How to read this report' for key to map.

Shelley Beach is located near the mouth of the Moruya River and backed by Eurobodalla National Park.

G

**Beach grade:** 


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream river sources.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain and often after 10 mm or more.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Nov 2020 to Mar 2024	91%	100	Stable	$\bigcirc$


#### **Sanitary inspection: Moderate**




#### Dry and wet weather water quality



#### **Microbial Assessment Category: B**





## **Tuross Main Beach**





See 'How to read this report' for key to map.

Tuross Main Beach is a 250 m long beach located between Tuross Headland in the north and Tuross Lake in the south.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but can be susceptible to pollution after rain, with several potential sources of faecal contamination including upstream river sources.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit in response to little or no rain and often after 10 mm or more.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Nov 2020 to Mar 2024	93%	100	Stable (	$\bigcirc$


### **Sanitary inspection: Low**




### Dry and wet weather water quality



### **Microbial Assessment Category: B**





## **Brou Beach**





See 'How to read this report' for key to map.

Brou Beach is located to the north of Dalmeny. The beach is approximately 6.5 km long and is backed by Eurobodalla National Park.


The Beach Suitability Grade of Very Good indicates microbial water quality is considered suitable for swimming almost all of the time, with few potential sources of significant faecal contamination.

Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit after 10 mm or more of rain.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Ocean beach	Nov 2020 to Mar 2024	98%	100	Stable	$\bigcirc$


### Sanitary inspection: Low




### Dry and wet weather water quality



### **Microbial Assessment Category: A**





## Wagonga Inlet





See 'How to read this report' for key to map.

The swimming site is a netted enclosure at the mouth of Wagonga Inlet. The town of Narooma is located on the southern side of the inlet.


The Beach Suitability Grade of Good indicates microbial water quality is considered suitable for swimming most of the time but may be susceptible to pollution after rain, with potential faecal contamination from upstream sources in Wagonga Inlet.

Enterococci levels increased slightly with increasing rainfall, occasionally exceeding the safe swimming limit after little or no rain, and often after 10 mm or more.


The site has been monitored since 2002.

Site	Assessment	Dry weather samples suitable for swimming	Water	Beach grade	
type	period		samples	status	
Estuarine	Nov 2020 to Mar 2024	95%	100	Stable (	$\bigcirc$


#### **Sanitary inspection: Moderate**




### Dry and wet weather water quality



### **Microbial Assessment Category: B**





## Narooma Main Beach





microbial water quality is considered suitable for swimming most of the time but may be

summer months.

susceptible to pollution after rain, with potential sources of faecal contamination including outflow from Little Lake.

Narooma Beach is approximately 750 m long and is patrolled on weekends and holidays during the


The Beach Suitability Grade of Good indicates

Enterococci levels increased slightly with increasing rainfall, often exceeding the safe swimming limit after 10 mm or more of rain.


See 'How to read this report' for key to map.

The site has been monitored since 2002. Site Assessment Dry weather samples Water **Beach grade** type period suitable for swimming samples status Ocean beach Nov 2020 to Stable 97% 100 Mar 2024

### Sanitary inspection: Low





### Dry and wet weather water quality



=40 cfu/100mL =41-200 cfu/100mL = 201-500 cfu/100mL =>500 cfu/100mL

### **Microbial Assessment Category: B**





## How to read this report

## **Beach Suitability Grades**

Beach Suitability Grades provide an assessment of the suitability of a swimming location for recreation over time and are based on a combination of sanitary inspection (identification and rating of potential pollution sources at a beach) and microbial assessment (water quality measurements gathered over previous years). There are 5 grades ranging from Very Good to Very Poor:



Location has generally excellent microbial water quality and very few potential sources of faecal pollution. Water is considered suitable for swimming almost all of the time

## G Good

Location has generally good microbial water quality and water is considered suitable for swimming most of the time. Swimming should be avoided during and for up to one day following heavy rain at ocean beaches and up to 3 days at estuarine sites



Microbial water quality is generally suitable for swimming, but because of the presence of significant sources of faecal contamination, extra care should be taken to avoid swimming during and for up to 3 days following rainfall or if there are signs of pollution such as discoloured water or odour or debris in the water Some Beach Suitability Grades in this report are **provisional**, as the information required for the analysis is incomplete due to limited bacterial data or limited information on potential pollution sources in the catchment.

## P Poor

Location is susceptible to faecal pollution and microbial water quality is not always suitable for swimming. During dry weather conditions, ensure that the swimming location is free of signs of pollution, such as discoloured water, odour or debris in the water, and avoid swimming at all times during and for up to 3 days following rainfall

## VP Very Poor

Location is very susceptible to faecal pollution and microbial water quality may often be unsuitable for swimming. It is generally recommended to avoid swimming at these sites almost all of the time.

## Follow Up

Sometimes a location's sanitary inspection and water quality data produce incongruent results. These locations are classified as 'Follow Up'. Further assessment will be required to obtain the necessary data to provide a definite classification in accordance with national guidelines.

### The guidelines

The National Health and Medical Research Council's guidelines for managing risks in recreational water (NHMRC 2008) were adopted for use in New South Wales in May 2009. These guidelines have been adopted in all Australian states and territories and are supported by guidance notes developed by the Department of Health Western Australia (WA Department of Health 2007).

## Enterococci

The national guidelines advocate the use of enterococci as the single preferred faecal indicator in recreational waters.

These bacteria are excreted in faeces and are rarely present in unpolluted waters. Enterococci have shown a clear dose–response relationship to disease outcomes in

marine waters in the northern hemisphere. In accordance with the guidelines, Beachwatch tests for enterococci only. The enterococci density in water samples is analysed in the laboratory using method AS/NZS 4276.9:2007 (Standards Australia 2007).

Enterococci are measured in colony forming units per 100 mL of sample (cfu/100 mL).

		Microbial Assessment Category			
		А	В	С	D
Sanitary Inspection Category	Very Low	Very Good	Very Good	Follow Up	Follow Up
	Low	Very Good	Good	Follow Up	Follow Up
	Moderate	Good	Good	Poor	Poor
	High	Good	Fair	Poor	Very Poor
	Very High	Follow Up	Fair	Poor	Very Poor

Beach Suitability Grades are determined by using the following matrix:

\* Follow up occurs when sanitary inspection and water quality data produce potentially incongruent results; further assessment will be required.

Using the Beach Suitability Grade classification matrix, sites assigned a moderate Sanitary Inspection Category can only be rated as Good or Poor, with no option of Fair grades. This can create the impression of a large change in water quality when in fact there need only be a slight increase in bacterial counts to push it over the threshold, with no significant increase in the risk to public health.

## Microbial Assessment Category (MAC)

There are 4 Microbial Assessment Categories (A to D) and these are determined from the 95th percentile of an enterococci dataset of at least 100 data points. Each MAC is associated with a risk of illness determined from epidemiological studies. The risks of illness shown below are not those associated with a single data point but are the overall risk of illness associated with an enterococci dataset with that 95th percentile (Wyer et al. 1999).

## Risk of illness associated with Microbial Assessment Categories

Category	<b>Enterococci</b> (cfu/100 mL)	Illness risk*
A	≤40	GI illness risk: <1% AFR illness risk: <0.3%
В	41–200	GI illness risk: 1–5% AFR illness risk: 0.3–1.9%
С	201-500	GI illness risk: >5–10% AFR illness risk: >1.9– 3.9%
D	>500	GI illness risk: >10% AFR illness risk: >3.9%

\* GI = gastrointestinal illness; AFR = acute fever and rash

### **Calculating the MAC**

The 95th percentile is a useful statistic for summarising the distribution of enterococci data at a site. It embodies elements of both the location of the distribution (how high/low the enterococci counts are) and the scale of the distribution (how variable the enterococci counts are).

The 95th percentile values for each of the 4 Microbial Assessment Categories were determined by the World Health Organization using enterococci data collected from swimming locations across Europe. These values will represent different probabilities of illness if the distribution of enterococci data from swimming locations in New South Wales differs from the European distribution.

In recognition of this issue, Dr Richard Lugg (Department of Health, Western Australia) has developed a Microsoft<sup>®</sup> Excel tool for calculating a modified 95th percentile that takes into account the distribution of data. The WA Department of Health recommends a minimum of 65 samples, collected from a particular site over 5 consecutive years, to provide sufficient confidence and reliability in the 95th percentile data output. This tool has been used to calculate the 95th percentile values

presented in this report and has been adopted for use by other state governments in Australia.

The tool can be downloaded from the WA Government's 'Environmental waters publications' webpage, under *Forms and templates*.

## Sanitary Inspection Category (SIC)

The aim of a sanitary inspection is to identify all sources of faecal contamination that could affect a swimming location and assess the risk to public health posed by these sources. It is an assessment of the likelihood of bacterial contamination from identified pollution sources and should, to some degree, correlate with the bacterial water quality results obtained from sampling.

The main sources of faecal contamination considered in the sanitary inspection are: bathers, toilet facilities, wastewater treatment plants (WWTPs), sewage overflows, sewer chokes, onsite systems, wastewater reuse, stormwater, river discharge, lagoons, boats and animals.

Rivers, lakes and estuaries themselves can be potential sources of faecal contamination to sites located in these waterbodies, with contaminated water from upstream or surrounding areas impacting water quality at the swimming location. This source is captured in river discharge or lagoon category, and shown as the waterbody in the sanitary inspection charts.

Through the sanitary inspection process, beaches are categorised to reflect the overall likelihood of faecal contamination. There are 5 categories: Very Low, Low, Moderate, High and Very High.

More information about the **sanitary inspection** process is available in the Beachwatch Protocol for assessment and management of microbial risks in recreational waters, found on the department's website.



Stormwater drain flow Photo: Beachwatch/DCCEEW

Stormwater in urban areas often contains sewage from leakages, overflows or sewer chokes when the sewerage system fails.

Sewage overflows can occur in wet weather when the network has exceeded capacity due to rainwater entering the system. The mix of sewage and rainwater discharges from designated overflow points and drains to waterways, usually via the stormwater system. Overflows from the sewerage system can also occur in dry weather due to mechanical failure or power outage.

Sewer chokes occur due to blockages in the pipes usually due to tree roots, oil, grease or debris. This causes sewage to back up and escape via sewer inspection points, designed overflow structures or cracks in the pipes, then drain to waterways, usually via the stormwater system.

## **Explanation of tables**

Each region contains tables listing all monitored swimming sites including site type, beach grade and change in grade from the previous year.

The following symbols are used to show the change in beach grade from the previous year:



Stable



Improved




Declined

A provisional grade indicates the assessment is based on limited data collected during the assessment period and should not be compared to the beach grade from the previous year.

# Explanation of graphs, charts, and information bars on beach pages


## Microbial Assessment Category (MAC) chart

On each beach page, the MACs for the last 5 years are displayed on a simple bar chart. The MAC for the current year is based on enterococci data collected during the assessment period. The bars are labelled with the 95th percentile value for each year and the thresholds dividing the A, B, C and D categories are marked in green, amber and red for reference.




## Sanitary Inspection Category (SIC) chart

The results of the sanitary inspection for each swimming location are presented in a radar pie chart. The chart shows the likelihood that each identified pollution source will contribute to faecal contamination at a swimming site, as indicated by the size and colour of the segment, ranging from very low (lightest colour) to very high (darkest colour) as shown below. The sum of these contributions is the overall likelihood, or Sanitary Inspection Category.



### Wet and dry weather water quality chart

Enterococci levels in wet and dry weather conditions are presented for each swimming location as a bar graph. All data collected during the assessment period is included in the analysis. Dry weather is defined as no rainfall recorded in the previous 24 hours. Each bar is colour coded to show the number of enterococci results up to 40 cfu/100 mL, between 41 and 200 cfu/100 mL, between 201 and 500 cfu/100 mL and greater than 500 cfu/100 mL. These categories reflect the Microbial Assessment Category thresholds and are coloured on the graph as dark green, light green, amber and red respectively.




■<=40 cfu/100mL ■41-200 cfu/100mL =201-500 cfu/100mL =>500 cfu/100mL

It is expected that swimming sites with lower levels of flushing will show some elevated bacterial results in dry weather samples (no rainfall in the previous 24 hours) due to the longer time needed to recover from a rainfall event. At some estuarine and lake/lagoon swimming locations the impacts of stormwater pollution on beach water quality may be detected up to 3 days after rainfall.


### Water quality in response to rainfall

Trends in enterococci levels in response to rainfall are shown using a box plot. For reference, enterococci levels of 40 cfu/100 mL and 200 cfu/100 mL are indicated with a green and orange line, respectively. The 40 cfu/100 mL level is referred to as the 'safe swimming limit'. The enterococci data were obtained from the last 5 years of monitoring. Rainfall data were obtained from rain gauges situated close to the sample site and are 24-hour totals to 9 am on the day of sampling. If there are fewer than 5 enterococci data points in a rainfall category, individual data points are presented instead of a box plot. At sites

where many results are below the detection limit (1 cfu/100 mL), only the upper portion of the box plots will be visible.



Each part of the box plot represents a significant percentile value of the sample population:



### Information bars

Information bars on each beach page provide a summary of details about the swimming site.

The **assessment period** shows the timeframe in which the water samples were collected. The NHMRC guidelines state beach grades should be determined from the most recent 100 water quality results collected within a 5-year period. The assessment period varies between sites depending on sampling frequency.

Dry weather samples suitable for swimming (**dry weather swimmability**) shows the percentage of water samples with enterococci levels below 40 cfu/100 mL. Dry weather is defined as no rainfall in the previous 24 hours. Swimming sites with lower levels of flushing often have a

lower percentage of dry weather samples within the safe swimming limit due to the impacts of rainfall detected up to 3 days after the event.

### **Explanation of maps**

A map of individual swimming locations is presented on each beach page. The scale of the maps is 1:10,000. Each map shows the location of the sampling site, land use and features such as surf lifesaving clubs. Potential pollution sources such as stormwater drains, sewage pumping stations, wastewater treatment plants, lagoons, rivers and creeks, are shown where accurate data is held.

Key to maps			
	Sampling Site		
	Surf Life Saving Club		
$\odot$	Wastewater Treatment Plant		
S PS	Sewage Pumping Station		
0	Sewage Overflow		
0	Stormwater Drain		
	Water		
	Baths		
	National Park/Reserve/ Other Park		
	Built-up Area		
	Sand		
	Roads		
	Major Roads		
-	Baths – Netted Area		
-	Breakwater/Wharf		

## References

NHMRC (2008) *Guidelines for managing risks in recreational water*, National Health and Medical Research Council, Australian Government Publishing Service, Canberra, ACT.

Standards Australia (2007) *AS/NZS* 4276.9:2007, *Water microbiology Method* 9: *Enterococci* – *Membrane filtration method* (*ISO* 7899-2:2000, *MOD*), Standards Australia International Ltd, Sydney and Standards New Zealand, Wellington.

WA Department of Health (2007), Microbial quality of recreational water guidance notes in support of chapter 5 of the National Health and Medical Research Council guidelines for managing risks in recreational water, 2006, Department of Health, Western Australia and The University of Western Australia, October 2007, ww2.health.wa.gov.au/Articles/A\_E/Environmentalwaters-publications, accessed 23/06/23.

Wyer MD, Kay D, Fleisher JM, Salmon RL, Jones F, Godfree AF, Jackson G and Rogers A (1999) 'An experimental health related classification for marine waters', *Water Research*, 33(3):715–722.

## More information

- Beachwatch webpage
- <u>Coastal management program progress</u>
- Sanitary inspection of beaches
- WA Government environmental water publications
- Shoalhaven Council Coast and waterways
- Eurobodalla Shire Council Coast and waterways